2 · Pure Mathematics 2

Pearson Edexcel · International A-Level · Mathematics · Vocabulary in context

Word	Example
laws of logarithms /lɔ:z ɒv 'lɒgərɪθəmz/	The **laws of logarithms** say ln(ab) = ln a + ln b. ln(ab) = ln a + ln b
trigonometric identities /ˌtrɪgənəʊ'metrɪk aɪ'dentɪtiz/	**Trigonometric identities** like $\sin^2 x + \cos^2 x = 1$ are always true. $\sin^2 x + \cos^2 x = 1$
proof /pru:f/	A **proof** must justify every case covered by its claim.
proof by exhaustion /pru:f baɪ eg'zɔ:stʃn/	In **proof by exhaustion**, we check every possible case one by one.
disproof by counterexample /dɪ'spru:f baɪ 'kaʊntəregzæmpl/	**Disproof by counterexample** shows a claim is false with one case.
factor theorem /'fæktə 'θɪərəm/	The **factor theorem** says f(a) = 0 exactly when (x-a) is a factor of f(x). f(a) = 0 (x-a) f(x)
remainder theorem /rɪ'meɪndə 'θɪərəm/	The **remainder theorem** says f(a) is the remainder when f(x) is divided by (x − a). f(x) (x − a) f(a)
algebraic division /ˌældʒɪ'breɪɪk dɪ'vɪʒn/	Use **algebraic division** to divide a polynomial by (x − 2). (x − 2)
quotient /'kwəʊʃənt/	When you divide 12 by 3, the **quotient** is 4. 12 3 4
chord /kɔ:d/	A **chord** is a straight line joining two points on a circle.

exercises.lol exercises.com.cn

Word	Example
tangent /'tændʒənt/	Draw a **tangent** to the curve to find the gradient at one point.
sequence /'si:kwəns/	A **sequence** places its terms in a particular order.
increasing /ɪn'kri:sɪŋ/	A sequence is **increasing** if each term is bigger than the last.
decreasing /'di:kri:sɪŋ/	A sequence is **decreasing** if each term is smaller than the last.
periodic /ˌpɪərɪ'ɒdɪk/	sin x is **periodic**: it repeats every 360°. sin x 360°
arithmetic sequences /ə'rɪθmətɪk 'si:kwənsɪz/	In **arithmetic sequences**, you add the same number each time.
geometric sequences /ˌdʒi:ə'metrɪk 'si:kwənsɪz/	In **geometric sequences**, you multiply by the same number each time.
binomial expansion /baɪ'nəʊmɪəl ek'spænʃn/	The **binomial expansion** writes $(1 + x)^n$ as a sum of terms. $(1 + x)^n$
logarithm /'lɒgərɪθəm/	Take the **logarithm** of both sides.
domain /də'meɪn/	Leave zero out of the **domain** when x is on the bottom of a fraction. x
trigonometric equations /ˌtrɪgənəʊ'metrɪk ɪ'kweɪʒnz/	Solve the **trigonometric equations** for $0° \leq x < 360°$. $0° \leq x < 360°$
stationary points /'steɪʃənəri pɔɪnts/	At **stationary points**, the gradient is zero.
maximum /'mæksɪməm/	A **maximum** point is the top of a hill on the graph.

Word	Example
minimum /ˈmɪnɪməm/	A **minimum** point is the bottom of a valley.
definite integral /ˈdefɪnət ˈɪntɪgrəl/	A **definite integral** has limits and gives a number, such as an area.
area under the curve /ˈeərɪə ˈʌndə ðə kɜːv/	Integration finds the **area under the curve**.
trapezium rule /trəˈpiːzɪəm ruːl/	The **trapezium rule** estimates an integral using straight-sided strips.

exercises.lol exercises.com.cn