

Exercise walk-through

3.5

Differentiation rules and implicit work

Pearson Edexcel · International A-Level · Mathematics

Check yourself

Check yourself

Rational simplification and inverses verified by composing; modulus cases checked; combined transformations; sec, cosec, cot with identities; harmonic form with exact R; compound and double angle proofs; log-graph recovery; growth-decay models with floor; product, quotient, chain and implicit differentiation; recognition integrals; sign-change and iteration methods.

Exercises 2.1·2.3

2.1 Differentiate $y = \sin 5x$. [1]

2.2 Differentiate $y = e^{-3x}$. [1]

2.3 Differentiate $y = x^2 \cos x$ using the product rule. [3]

Exercises 2.1·2.3

2.1 Differentiate $y = \sin 5x$. [1]

2.1 $\frac{dy}{dx} = 5 \cos 5x$.

2.2 Differentiate $y = e^{-3x}$. [1]

2.2 $\frac{dy}{dx} = -3e^{-3x}$.

2.3 Differentiate $y = x^2 \cos x$ using the product rule. [3]

2.3 $\frac{dy}{dx} = 2x \cos x - x^2 \sin x$.

Exercises 2.4·2.6

2.4 Differentiate $y = \frac{e^x}{x}$ using the quotient rule. [3]

2.5 Differentiate $y = \ln(4x^3 + 1)$ using the chain rule. [2]

2.6 Find $\frac{dy}{dx}$ for $x = e^{2y}$. [3]

Exercises 2.4·2.6

2.4 Differentiate $y = \frac{e^x}{x}$ using the quotient rule. [3]

$$\mathbf{2.4} \quad \frac{dy}{dx} = \frac{e^x \cdot x - e^x \cdot 1}{x^2} = \frac{e^x(x-1)}{x^2}.$$

2.5 Differentiate $y = \ln(4x^3 + 1)$ using the chain rule. [2]

$$\mathbf{2.5} \quad \frac{dy}{dx} = \frac{12x^2}{4x^3 + 1} \text{ (chain: outer } \ln, \text{ inner } 4x^3 + 1).$$

2.6 Find $\frac{dy}{dx}$ for $x = e^{2y}$. [3]

$$\mathbf{2.6} \quad \frac{dx}{dy} = 2e^{2y} = 2x, \text{ so } \frac{dy}{dx} = \frac{1}{2x}.$$

Exercise 3.1

3.1 (*Adapted from Edexcel IAL WMA13 June 2025 Q7.*) The curve C has equation $y = e^{-x^2} \sin 3x$ for $0 \leq x \leq 3$. Show, using calculus, that the x -coordinates of the stationary points of C satisfy

$$x = \frac{1}{3} \arctan \frac{3}{2x}$$

[4]

Exercise 3.1

3.1 (*Adapted from Edexcel IAL WMA13 June 2025 Q7.*) The curve C has equation $y = e^{-x^2} \sin 3x$ for $0 \leq x \leq 3$. Show, using calculus, that the x -coordinates of the stationary points of C satisfy

$$x = \frac{1}{3} \arctan \frac{3}{2x}$$

[4]

3.1 From the worked derivative: stationary points have

$$e^{-x^2} (3 \cos 3x - 2x \sin 3x) = 0$$

$e^{-x^2} > 0$ always, so $3 \cos 3x = 2x \sin 3x$. Dividing by $\cos 3x$ (at a stationary point $\cos 3x \neq 0$, since that would force $\sin 3x = \pm 1 \neq 0$ and then $0 = \pm 2x$ fails for $x > 0$):

$$3 = 2x \tan 3x \implies \tan 3x = \frac{3}{2x} \implies 3x = \arctan \frac{3}{2x} \implies x = \frac{1}{3} \arctan \frac{3}{2x} \quad \blacksquare$$

Exercise 3.2

3.2 A cylindrical tank drains so that the volume of water is $V = 500e^{-0.1t}$ litres after t minutes. Find the rate at which water leaves at $t = 10$, and the time at which the rate is 5 litres per minute. [5]

Exercise 3.2

3.2 A cylindrical tank drains so that the volume of water is $V = 500e^{-0.1t}$ litres after t minutes. Find the rate at which water leaves at $t = 10$, and the time at which the rate is 5 litres per minute. [5]

3.2 $\frac{dV}{dt} = -50e^{-0.1t}$; at $t = 10$: $-50e^{-1} = -18.4$ litres per minute (leaving at 18.4 L/min). Rate = 5: $50e^{-0.1t} = 5 \Rightarrow e^{-0.1t} = 0.1 \Rightarrow t = 10 \ln 10 = 23.0$ minutes (3 s.f.). Check: $V(23.0) = 500e^{-2.303} = 50.0$ litres, one tenth of the initial – matching the rate being one tenth of the initial 50 L/min ✓.

Local review guidance: correct rule with both parts differentiated; simplified derivative; exponential-rate equation solved with logs. This is authored classroom guidance.