

Exercise walk-through

4.4

Implicit differentiation and differential equations

Pearson Edexcel · International A-Level · Mathematics

Check yourself

Check yourself

Contradiction proofs with the assumption named; partial fractions with repeated factors and improper cases; parametric conversions and gradients; the general binomial series with validity; implicit tangents; separable equations with conditions; parts, substitution and partial-fraction integrals; exact volumes of revolution; vector angles from the scalar product.

Exercises 2.1·2.3

2.1 Find $\frac{dy}{dx}$ for $x^2 + y^2 = 25$. [2]

2.2 Find $\frac{dy}{dx}$ for $x^2 + 2xy = 3$. [3]

2.3 A cube's edge grows at 0.1 cm/s. Find $\frac{dV}{dt}$ when the edge is 5 cm. [3]

Exercises 2.1·2.3

2.1 Find $\frac{dy}{dx}$ for $x^2 + y^2 = 25$. [2]

2.1 $2x + 2y\frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{x}{y}$.

2.2 Find $\frac{dy}{dx}$ for $x^2 + 2xy = 3$. [3]

2.2 $2x + 2y + 2x\frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{x+y}{x}$.

2.3 A cube's edge grows at 0.1 cm/s. Find $\frac{dV}{dt}$ when the edge is 5 cm. [3]

2.3 $V = x^3: \frac{dV}{dt} = 3x^2\frac{dx}{dt} = 3(25)(0.1) = 7.5 \text{ cm}^3/\text{s}.$

Exercises 2.4·3.1

2.4 Solve $\frac{dy}{dx} = \frac{2x}{y}$ with $y = 3$ at $x = 0$. [4]

3.1 (*Adapted from Edexcel IAL WMA14 June 2025 Q2.*) A sealed spherical balloon loses air at a constant 12 cm^3 per second. Show that the rate of decrease of its radius is inversely proportional to the square of the radius, and find that rate when $r = 3 \text{ cm}$. [5]

Exercises 2.4·3.1

2.4 Solve $\frac{dy}{dx} = \frac{2x}{y}$ with $y = 3$ at $x = 0$. [4]

2.4 $\int y \, dy = \int 2x \, dx \Rightarrow \frac{y^2}{2} = x^2 + c$; $y(0) = 3$ gives $c = \frac{9}{2}$, so $y^2 = 2x^2 + 9$.

3.1 (*Adapted from Edexcel IAL WMA14 June 2025 Q2.*) A sealed spherical balloon loses air at a constant 12 cm^3 per second. Show that the rate of decrease of its radius is inversely proportional to the square of the radius, and find that rate when $r = 3 \text{ cm}$. [5]

3.1 $V = \frac{4}{3}\pi r^3$, $\frac{dV}{dr} = 4\pi r^2$; chain rule with $\frac{dV}{dt} = -12$:

$$-12 = 4\pi r^2 \frac{dr}{dt} \implies \frac{dr}{dt} = -\frac{3}{\pi r^2}$$

The rate of decrease $\frac{3}{\pi r^2}$ is a constant multiple of $\frac{1}{r^2}$ – inversely proportional to r^2 . ■

At $r = 3$:

$$\frac{3}{9\pi} = \frac{1}{3\pi} \approx 0.106 \text{ cm/s (3 s.f.)}$$

Check: units $\text{cm}^3/\text{s} \div \text{cm}^2 = \text{cm/s} \checkmark$.

Exercise 3.2

3.2 (*Adapted from Edexcel IAL WMA14 June 2025 Q3.*) Solve the differential equation

$$y \frac{dy}{dx} = 6 \sec 3x \tan 3x, \quad y = 2 \text{ at } x = 0$$

giving your answer in the form $y^2 = g(x)$. [6]

Exercise 3.2

3.2 (*Adapted from Edexcel IAL WMA14 June 2025 Q3.*) Solve the differential equation

$$y \frac{dy}{dx} = 6 \sec 3x \tan 3x, \quad y = 2 \text{ at } x = 0$$

giving your answer in the form $y^2 = g(x)$. [6]

3.2 Separate: $\int y \, dy = \int 6 \sec 3x \tan 3x \, dx$ with $\int \sec 3x \tan 3x \, dx = \frac{1}{3} \sec 3x$:

$$\frac{y^2}{2} = 2 \sec 3x + c \implies y^2 = 4 \sec 3x + C$$

Condition $y = 2$ at $x = 0$: $4 = 4(1) + C \Rightarrow C = 0$:

$$y^2 = 4 \sec 3x$$

Check: $\sqrt{4} = 2$ at $x = 0$ ✓; differentiating back: $2y \frac{dy}{dx} = 12 \sec 3x \tan 3x \Rightarrow y \frac{dy}{dx} = 6 \sec 3x \tan 3x$ ✓.

Local review guidance: correct implicit/rate structure; exact derivative or particular solution; condition applied and the solution checked. This is authored classroom guidance.