

Exercise walk-through

5.5

Momentum and impulse

Pearson Edexcel · International A-Level · Mathematics

Check yourself

Check yourself

State modelling assumptions and their consequences; vector velocities, positions, bearings; multi-phase suvat and motion graphs; paired equations over a pulley with the 2T pulley force; impulse on one body and momentum conservation with rebounds; μR moving versus capped at rest; equilibrium by two resolutions; moments about successive tilting pivots.

Exercises 2.1·2.3

2.1 A ball of mass 0.5 kg hits a wall perpendicular to it at 6 m s^{-1} and rebounds at 4 m s^{-1} . Find the magnitude of the impulse the wall exerts on the ball. [3]

2.2 A 2 kg particle moving at 5 m s^{-1} collides with a 3 kg particle at rest and they move off together. (a) Find their common speed. (b) Find the impulse on the 3 kg particle. [4]

2.3 A ball of mass 0.4 kg is dropped and hits the floor at 5 m s^{-1} , rebounding at 3 m s^{-1} . Find the impulse exerted by the floor. [4]

Exercises 2.1·2.3

2.1 A ball of mass 0.5 kg hits a wall perpendicular to it at 6 m s^{-1} and rebounds at 4 m s^{-1} . Find the magnitude of the impulse the wall exerts on the ball. [3]

2.1 Reversal: $I = 0.5(6 + 4) = 5 \text{ N s}$.

2.2 A 2 kg particle moving at 5 m s^{-1} collides with a 3 kg particle at rest and they move off together. (a) Find their common speed. (b) Find the impulse on the 3 kg particle. [4]

2.2 (a) $2(5) = (5)v$, so $v = 2 \text{ m s}^{-1}$. (b) Impulse on the 3 kg: $3(2 - 0) = 6 \text{ N s}$.

2.3 A ball of mass 0.4 kg is dropped and hits the floor at 5 m s^{-1} , rebounding at 3 m s^{-1} . Find the impulse exerted by the floor. [4]

2.3 Reversal: $I = 0.4(5 + 3) = 3.2 \text{ N s}$.

Exercises 2.4·3.1

2.4 A 1 kg particle moving right at 4 m s^{-1} collides with a 2 kg particle moving left at 1 m s^{-1} . After the collision the 1 kg particle moves left at 0.5 m s^{-1} . Find the velocity of the 2 kg particle. [4]

3.1 (*Edexcel IAL WME01 June 2025 Q1.*) Two particles P and Q move in the same direction along the same straight line on a smooth horizontal surface. The mass of P is km and the mass of Q is $3m$. They collide directly. Immediately before the collision, the speed of P is $5u$ and the speed of Q is $2u$. Immediately after, the speed of P is $2u$ with direction unchanged, and the speed of Q is v . The impulse exerted on Q by P has magnitude $4.5mu$. (a) Find v in terms of u . (b) Find k . [6]

Exercises 2.4·3.1

2.4 A 1 kg particle moving right at 4 m s^{-1} collides with a 2 kg particle moving left at 1 m s^{-1} . After the collision the 1 kg particle moves left at 0.5 m s^{-1} . Find the velocity of the 2 kg particle. [4]

2.4 Right positive: $1(4) + 2(-1) = 1(-0.5) + 2v$, so $2 = -0.5 + 2v$ and $v = 1.25 \text{ m s}^{-1}$ to the right.

3.1 (*Edexcel IAL WME01 June 2025 Q1.*) Two particles P and Q move in the same direction along the same straight line on a smooth horizontal surface. The mass of P is km and the mass of Q is $3m$. They collide directly. Immediately before the collision, the speed of P is $5u$ and the speed of Q is $2u$. Immediately after, the speed of P is $2u$ with direction unchanged, and the speed of Q is v . The impulse exerted on Q by P has magnitude $4.5mu$. (a) Find v in terms of u . (b) Find k . [6]

3.1 (a) Impulse-momentum on Q: $3m(v - 2u) = 4.5mu$, so $v - 2u = 1.5u$ and $v = 3.5u$. (b) The impulse on P is equal in magnitude and opposite in effect (it slows P): $km(5u - 2u) = 4.5mu$, so $3k = 4.5$ and $k = 1.5$.

Exercise 3.2

3.2 (*Edexcel IAL WME01 June 2025 Q8d, condensed.*) A ball of mass 1.5 kg falls and hits the ground with speed 3.13 m s^{-1} , rebounding with speed 0.4 m s^{-1} . Find the magnitude of the impulse it receives. [4]

Exercise 3.2

3.2 (*Edexcel IAL WME01 June 2025 Q8d, condensed.*) A ball of mass 1.5 kg falls and hits the ground with speed 3.13 m s^{-1} , rebounding with speed 0.4 m s^{-1} . Find the magnitude of the impulse it receives. [4]

3.2 $I = 1.5(3.13 + 0.4) = 5.3 \text{ N s (5.30)}.$

Local review guidance: impulse-momentum applied to ONE named body; momentum conserved with all velocities signed consistently; reversals handled as sums of speeds in magnitude-form or clean sign algebra. This is authored classroom guidance.