

Exercise walk-through

8.4

Populations, samples and sampling distributions

Pearson Edexcel · International A-Level · Mathematics

Check yourself

Check yourself

State distribution assumptions in context; solve moment pairs; scale Poisson rates; normalise piecewise densities; medians, modes and skewness; uniform facts both directions; census versus sample; statistics; small sampling distributions; one- and two-tailed tests with critical regions and actual levels; continuity-corrected normal thresholds.

Exercises 2.1·2.3

2.1 State one advantage and one disadvantage of a sample survey compared with a census. [2]

2.2 For samples (X_1, X_2) from a population with mean μ , state which of $X_1 + 2X_2$, $\bar{X} - \mu$ and $3\bar{X}$ are statistics, with reasons. [3]

2.3 A fair coin is tossed twice; S = number of heads. Write out the sampling distribution of S . [2]

Exercises 2.1·2.3

2.1 State one advantage and one disadvantage of a sample survey compared with a census. [2]

2.1 Advantage: cheaper and faster than a census. Disadvantage: the results contain sampling error – they do not describe every member of the population.

2.2 For samples (X_1, X_2) from a population with mean μ , state which of $X_1 + 2X_2$, $\bar{X} - \mu$ and $3\bar{X}$ are statistics, with reasons. [3]

2.2 $X_1 + 2X_2$ and $3\bar{X}$ involve sample values only: statistics. $\bar{X} - \mu$ contains the unknown population parameter μ : not a statistic.

2.3 A fair coin is tossed twice; S = number of heads. Write out the sampling distribution of S . [2]

2.3 $P(S = 0) = \frac{1}{4}$, $P(S = 1) = \frac{1}{2}$, $P(S = 2) = \frac{1}{4}$.

Exercises 3.1·3.2

3.1 (*Edexcel IAL WST02 June 2025 Q4.*) A bag contains a large number of beads of which the ratio of black to white beads is 2:5. A player takes 3 beads at random from the bag and scores 4 points for each black bead and 1 point for each white bead. The random variable X is the total number of points scored. (a) Find the sampling distribution of X . (b) In a game a player takes n sets of 3 beads (each set replaced before the next); Y is the number of sets scoring exactly 9 points. Find the least n such that $P(Y \geq 1) > 0.95$. [9]

3.2 (*Original.*) Two fair dice are rolled; M is the larger score shown. (a) Find the sampling distribution of M . (b) Find $E(M)$. [7]

Exercises 3.1·3.2

3.1 (*Edexcel IAL WST02 June 2025 Q4.*) A bag contains a large number of beads of which the ratio of black to white beads is 2:5. A player takes 3 beads at random from the bag and scores 4 points for each black bead and 1 point for each white bead. The random variable X is the total number of points scored. (a) Find the sampling distribution of X . (b) In a game a player takes n sets of 3 beads (each set replaced before the next); Y is the number of sets scoring exactly 9 points. Find the least n such that $P(Y \geq 1) > 0.95$. [9]

3.1 (a) With $p_b = \frac{2}{7}$ (large bag: constant probability), the number of black beads is $B(3, \frac{2}{7})$ and $X = 3 + 3 \cdot (\text{blacks})$: $P(X = 3) = (\frac{5}{7})^3 = \frac{125}{343}$; $P(X = 6) = \binom{3}{1} \frac{2}{7} (\frac{5}{7})^2 = \frac{150}{343}$; $P(X = 9) = \binom{3}{2} (\frac{2}{7})^2 \frac{5}{7} = \frac{60}{343}$; $P(X = 12) = (\frac{2}{7})^3 = \frac{8}{343}$ (total $\frac{343}{343}$ ✓). (b) $Y \sim B(n, \frac{60}{343})$: $P(Y \geq 1) = 1 - (\frac{283}{343})^n > 0.95$, so $(\frac{283}{343})^n < 0.05$, $n \ln \frac{283}{343} < \ln 0.05$, $n > 15.6$: least $n = 16$.

3.2 (*Original.*) Two fair dice are rolled; M is the larger score shown. (a) Find the sampling distribution of M . (b) Find $E(M)$. [7]

3.2 (a) $P(M = m) = P(\max = m) = \binom{m}{6}^2 - \binom{m-1}{6}^2 = \frac{2m-1}{36}$ for $m = 1..6$: $\frac{1}{36}, \frac{3}{36}, \frac{5}{36}, \frac{7}{36}, \frac{9}{36}$, (sums to 1 ✓). (b) $E(M) = \sum m \cdot \frac{2m-1}{36}$: summing $m(2m-1) = 2m^2 - m$ over $m = 1..6$ gives $2(91) - 21 = 161$, so $E(M) = \frac{161}{36} = 4.47$.

Local review guidance: enumeration with the correct per-draw probability and binomial weights; distribution sums to 1; follow-up counts wrapped in $B(n, p)$ with the complement taken before logs. This is authored classroom guidance.