

1.1 Motion graphs and uniform acceleration

Name: _____ Class: _____ Date: _____ Total: 24 marks

KEY WORDS displacement • velocity • acceleration • gradient
• tangent • uniform acceleration • systematic error • light gate

Objective

Build the skills to answer WPH11 questions on **motion graphs** and **uniformly accelerated motion**: read gradients and areas, choose the right suvat equation, and finish every “deduce” question with a compared conclusion. This sheet covers Unit 1 statements 1–3 and core practical 1.

You must be able to:


- read acceleration from the gradient of a velocity–time graph and displacement from its area
- tell distance from displacement when areas have both signs
- draw and use a tangent on a curved graph, and estimate an area by counting squares
- choose between the four suvat equations by the quantities given
- finish *deduce whether...* questions with a comparison **and** a conclusion
- describe how to determine the acceleration of free fall, including the s – t^2 graph

1 Key knowledge and worked examples

Study these first. Each one shows a **method** that a question later on this sheet needs.

■ Gradient and area

On a velocity–time graph the **gradient** is the acceleration and the **area** between the line and the time axis is the displacement. Area below the axis is negative displacement; the distance adds the sizes of all the areas.



Worked example. Find the maximum acceleration shown on the graph.

- Known: the steepest straight part runs from (4 s, 1.8 m s^{−1}) to (16 s, 5.0 m s^{−1}). Why: acceleration is the gradient of a velocity–time graph.

$$a = \frac{\Delta v}{\Delta t}$$

$$a = \frac{5.0 \text{ m s}^{-1} - 1.8 \text{ m s}^{-1}}{16 \text{ s} - 4 \text{ s}} = 0.27 \text{ m s}^{-2}$$

- Check: read both values from the line itself, not from grid lines nearby; the unit is m s^{−2}.

■ Choosing a suvat equation

The four equations connect s , u , v , a , t for **constant** acceleration:

$$v = u + at \quad s = \frac{(u + v)t}{2} \quad s = ut + \frac{1}{2}at^2 \quad v^2 = u^2 + 2as$$

given	wanted	use
u, a, t	v	$v = u + at$
u, v, t	s	$s = \frac{(u+v)t}{2}$
u, a, t	s	$s = ut + \frac{1}{2}at^2$
u, v, a	s (no time!)	$v^2 = u^2 + 2as$

Worked example (deduce-and-compare). A ball leaves a bat at 12 m s^{−1} upwards. Deduce whether it rises more than 6.0 m. Take upwards as positive.

- Known: $u = +12 \text{ m s}^{-1}$, $v = 0$ at the top, $a = -9.81 \text{ m s}^{-2}$. No time, so use $v^2 = u^2 + 2as$.

$$v^2 = u^2 + 2as \Rightarrow s = \frac{v^2 - u^2}{2a}$$


$$s = \frac{0 - (12 \text{ m s}^{-1})^2}{2 \times (-9.81 \text{ m s}^{-2})} = \frac{-144}{-19.62} \text{ m} = 7.3 \text{ m}$$

- $7.3 \text{ m} > 6.0 \text{ m}$, so the ball **does** rise more than 6.0 m . The comparison and the conclusion earn the last mark.

Core practical 1: the acceleration of free fall

Drop a ball through two light gates (or film it against a ruler): measure the fall distance s and the time t , or the velocities at the gates. Then:


- one pair of values: $s = ut + \frac{1}{2}at^2$ with $u = 0$ gives $g = \frac{2s}{t^2}$;
- better: several fall distances, plotted as s against t^2 . Since $s = \frac{1}{2}gt^2$, the graph is a straight line through the origin with gradient $g/2$, so $g = 2 \times \text{gradient}$.



A straight line that misses the origin shows a **systematic error**, for example a constant distance offset. A timing delay can distort the relationship rather than merely changing its intercept. Repeating drops reduces random scatter; it cannot remove an offset. In a strobe photograph, a shorter flash means the ball travels less distance during the flash. The image is sharper and the position reading has less uncertainty, if the scale and image resolution are unchanged.

Curves: tangents and squares

When a velocity–time graph is curved, the acceleration **changes**: find the acceleration at one instant from the gradient of a tangent, and estimate the area by counting grid squares (count a square when at least half of it lies under the curve).



Worked example. Estimate the acceleration of the cyclist at $t = 2.0 \text{ s}$.

- Known: the curve; a tangent at $t = 2.0 \text{ s}$ passes through about $(0, 10.8)$ and $(4.0 \text{ s}, 4.4 \text{ m s}^{-1})$. Why: the tangent's gradient is the instantaneous acceleration.

$$a = \frac{\Delta v}{\Delta t} = \frac{4.4 \text{ m s}^{-1} - 10.8 \text{ m s}^{-1}}{4.0 \text{ s} - 0} = -1.6 \text{ m s}^{-2}$$


- Check: the curve is steepest at the start, so the tangent gradient here (1.6 m s^{-2} in size) should be larger than the average gradient over the whole 8 s (1.3 m s^{-2}) – it is.

2 Practice

Now use the methods above.

2.1 State what is represented by (a) the gradient of a velocity–time graph; (b) the area between the line and the time axis on a velocity–time graph. [2]

2.2 The velocity–time graph shows a lift that goes up, comes back down, and passes its starting floor. Upwards is positive.



(a) Calculate the acceleration of the lift between $t = 0$ and $t = 2.0$ s. [1]

(b) Calculate the displacement of the lift between $t = 6.0$ s and $t = 12.0$ s. [2]

(c) Calculate the total distance travelled by the lift in the 14 s. (*Hint: for distance, every area counts as positive.*) [2]

2.3 A ball is dropped from rest and hits the ground after 0.55 s.

(a) Use $s = ut + \frac{1}{2}at^2$ to show that the drop height is about 1.5 m. [2]

(b) State one reason why a measured time from a handheld stopwatch could make the calculated g too small, and one improvement. [2]

2.4 A car accelerates uniformly from 8.0 m s^{-1} to 20 m s^{-1} over 96 m. Calculate its acceleration. [3]

3 Exam-style questions

3.1 An object moves with constant non-zero acceleration. Which row describes its displacement–time graph and its velocity–time graph? [1]

- ☐ **A** both graphs are straight lines
- ☐ **B** the displacement–time graph is curved; the velocity–time graph is a straight sloped line
- ☐ **C** both graphs are curved
- ☐ **D** the displacement–time graph is a straight sloped line; the velocity–time graph is curved

3.2 A train's velocity–time graph is a straight line from 3.0 m s^{-1} at $t = 0$ to 9.0 m s^{-1} at $t = 4.0$ s. What is the displacement in those 4.0 s? [1]

- ☐ **A** 12 m
- ☐ **B** 24 m
- ☐ **C** 36 m
- ☐ **D** 48 m

3.3 In an experiment to determine g , a student plots s on the y -axis for a falling object.

(a) State what should be plotted on the x -axis to give a straight line through the origin, and how the gradient gives g . [2]

(b) The student's s against t^2 line has a small fixed vertical intercept but the expected gradient. Suggest a distance-measurement error that could cause this and state the effect on g calculated from the gradient. [2]

3.4 A skydiver leaves an aeroplane with zero vertical velocity. In the first 1.4 s she falls 9.2 m. Air resistance can be ignored if her mean acceleration is greater than $0.95g$. Deduce whether air resistance can be ignored. [4]

Solutions

2.1 (a) acceleration (b) displacement. *Distance needs the sizes of the areas; displacement uses their signs.*

2.2 (a) $a = \frac{\Delta v}{\Delta t} = \frac{2.0 \text{ m s}^{-1} - 0}{2.0 \text{ s}} = 1.0 \text{ m s}^{-2}$. (b) Areas: $t = 6\text{--}8 \text{ s}$: $\frac{1}{2} \times 2.0 \text{ s} \times 2.0 \text{ m s}^{-1} = +2.0 \text{ m}$; $t = 8\text{--}10 \text{ s}$: $\frac{1}{2} \times 2.0 \text{ s} \times (-2.0 \text{ m s}^{-1}) = -2.0 \text{ m}$; $t = 10\text{--}12 \text{ s}$: $2.0 \text{ s} \times (-2.0 \text{ m s}^{-1}) = -4.0 \text{ m}$. Displacement = $2.0 - 2.0 - 4.0 = -4.0 \text{ m}$. (c) Distance = $|+2.0| + |-2.0| + |-4.0|$ plus the $0\text{--}6 \text{ s}$ area $\frac{1}{2}(4.0 + 6.0) \text{ s} \times 2.0 \text{ m s}^{-1} = 10 \text{ m}$ and the $12\text{--}14 \text{ s}$ triangle $\frac{1}{2} \times 2.0 \text{ s} \times 2.0 \text{ m s}^{-1} = 2.0 \text{ m}$: total $10 + 2.0 + 2.0 + 4.0 + 2.0 = 20 \text{ m}$.

2.3 (a) $s = ut + \frac{1}{2}at^2 = 0 + \frac{1}{2} \times 9.81 \text{ m s}^{-2} \times (0.55 \text{ s})^2 = 1.5 \text{ m}$. (b) Reaction time makes t too long, so $g = 2s/t^2$ comes out too small; use light gates or video with a known frame rate instead.

2.4 No time is given, so $v^2 = u^2 + 2as$. $a = \frac{v^2 - u^2}{2s} = \frac{(20 \text{ m s}^{-1})^2 - (8.0 \text{ m s}^{-1})^2}{2 \times 96 \text{ m}} = \frac{336}{192} \text{ m s}^{-2} = 1.8 \text{ m s}^{-2}$.

3.1 B. Constant acceleration makes velocity change linearly (straight sloped velocity–time line) while displacement follows $s = ut + \frac{1}{2}at^2$ (curved). A is wrong because a straight displacement–time line means constant velocity; C and D mismatch the pair.

3.2 B. The area is a trapezium: $s = \frac{(u+v)}{2}t = \frac{3.0+9.0}{2} \text{ m s}^{-1} \times 4.0 \text{ s} = 24 \text{ m}$. A uses only u ; C multiplies $v \times t$; D doubles correctly counted halves.

3.3 (a) Plot t^2 on the x -axis; $s = \frac{1}{2}gt^2$, so the gradient equals $g/2$ and $g = 2 \times \text{gradient}$. (b) A fixed offset in the distance reading adds the same amount to every s ; it changes the vertical intercept but not the gradient, so $g = 2 \times \text{gradient}$ is unchanged.

3.4 $s = ut + \frac{1}{2}at^2$ with $u = 0$: $a = \frac{2s}{t^2} = \frac{2 \times 9.2 \text{ m}}{(1.4 \text{ s})^2} = 9.4 \text{ m s}^{-2}$. Compare: $\frac{9.4}{9.81} = 0.96 > 0.95$; since the calculated acceleration exceeds $0.95g$, air resistance **can** be ignored.