

Exercise walk-through

4.2

Parametric equations and differentiation

Pearson Edexcel · International A-Level · Pure Mathematics

Check yourself

Check yourself

Contradiction proofs with the assumption named; partial fractions with repeated factors and improper cases; parametric conversions and gradients; the general binomial series with validity; implicit tangents; separable equations with conditions; parts, substitution and partial-fraction integrals; exact volumes of revolution; vector angles from the scalar product.

Exercises 2.1·2.3

2.1 Convert $x = 3t$, $y = t^2$ to a Cartesian equation. [2]

2.2 For $x = t^2$, $y = 2t$ ($t > 0$), find $\frac{dy}{dx}$ at $t = 3$. [3]

2.3 For $x = \cos t$, $y = \sin t$, find $\frac{dy}{dx}$ and state the curve. [3]

Exercises 2.1·2.3

2.1 Convert $x = 3t$, $y = t^2$ to a Cartesian equation. [2]

2.1 $t = \frac{x}{3} \Rightarrow y = \frac{x^2}{9}$.

2.2 For $x = t^2$, $y = 2t$ ($t > 0$), find $\frac{dy}{dx}$ at $t = 3$. [3]

2.2 $\frac{dx}{dt} = 2t = 6$, $\frac{dy}{dt} = 2$: $\frac{dy}{dx} = \frac{2}{6} = \frac{1}{3}$.

2.3 For $x = \cos t$, $y = \sin t$, find $\frac{dy}{dx}$ and state the curve. [3]

2.3 $\frac{dy}{dx} = \frac{\cos t}{-\sin t} = -\cot t$; eliminating t gives $x^2 + y^2 = 1$ (the unit circle).

Exercises 2.4·3.1

2.4 A curve has $x = t + 1$, $y = t^2 - 3t$. Find the equation of the tangent at $t = 2$. [4]

3.1 (*Demand adapted from Edexcel IAL WMA14 June 2025.*) A curve is given by $x = 4t$,
 $y = \frac{4}{t}$ ($t > 0$).

(a) Find $\frac{dy}{dx}$ in terms of t .

(b) Find the equation of the normal at $t = 2$, in the form $ax + by + c = 0$ with integer coefficients. [6]

Exercises 2.4·3.1

2.4 A curve has $x = t + 1$, $y = t^2 - 3t$. Find the equation of the tangent at $t = 2$. [4]

2.4 $\frac{dy}{dx} = \frac{2t-3}{1}$; at $t = 2$: gradient 1, point $(3, -2)$. Tangent: $y + 2 = x - 3 \implies y = x - 5$.

3.1 (*Demand adapted from Edexcel IAL WMA14 June 2025.*) A curve is given by $x = 4t$, $y = \frac{4}{t}$ ($t > 0$).

(a) Find $\frac{dy}{dx}$ in terms of t .

(b) Find the equation of the normal at $t = 2$, in the form $ax + by + c = 0$ with integer coefficients. [6]

3.1 (a) $\frac{dx}{dt} = 4$, $\frac{dy}{dt} = -\frac{4}{t^2}$:

$$\frac{dy}{dx} = -\frac{1}{t^2}$$

(b) At $t = 2$: gradient $-\frac{1}{4}$, point $(8, 2)$. Normal gradient 4: $y - 2 = 4(x - 8) \implies 4x - y - 30 = 0$. Check: $4(8) - 2 - 30 = 0 \checkmark$; the gradients multiply to $-\frac{1}{4} \times 4 = -1 \checkmark$.

Exercise 3.2

3.2 A curve has $x = t^2$, $y = t^3$. Find the Cartesian equation and the tangent at $t = -1$. [6]

Exercise 3.2

3.2 A curve has $x = t^2$, $y = t^3$. Find the Cartesian equation and the tangent at $t = -1$. [6]

3.2 Cartesian: $t^2 = x$, $t^3 = y \Rightarrow y^2 = t^6 = (t^2)^3 = x^3$, so $y^2 = x^3$. At $t = -1$: point $(1, -1)$, $\frac{dy}{dx} = \frac{3t^2}{2t} = \frac{3t}{2} = -\frac{3}{2}$. Tangent: $y + 1 = -\frac{3}{2}(x - 1) \implies 3x + 2y - 1 = 0$. Check: $3(1) + 2(-1) - 1 = 0$ ✓; and $(1, -1)$ satisfies $y^2 = x^3$ ✓.

Local review guidance: both derivatives with respect to t found; gradient value at the parameter; line through the correct point in the requested form. This is authored classroom guidance.