

Exercise walk-through

5.3

Roots of quadratic equations

Pearson Edexcel · International A-Level · Pure Mathematics

Exercises 2.1·2.3

2.1 For $2x^2 - 8x + 3 = 0$ with roots α, β , find (a) $\alpha + \beta$; (b) $\alpha\beta$; (c) $\alpha^2 + \beta^2$; (d) $\frac{1}{\alpha} + \frac{1}{\beta}$. [4]

2.2 The equation $x^2 + 6x + 4 = 0$ has roots α, β . Without solving it, form a quadratic equation with roots 3α and 3β . [3]

2.3 For $x^2 - 5x + 9 = 0$ with roots α, β , find the value of $\frac{\alpha}{\beta} + \frac{\beta}{\alpha}$. [3]

Exercises 2.1·2.3

2.1 For $2x^2 - 8x + 3 = 0$ with roots α, β , find (a) $\alpha + \beta$; (b) $\alpha\beta$; (c) $\alpha^2 + \beta^2$; (d) $\frac{1}{\alpha} + \frac{1}{\beta}$. [4]

2.1 (a) $\alpha + \beta = 4$; (b) $\alpha\beta = \frac{3}{2}$; (c) $16 - 3 = 13$; (d) $\frac{4}{3/2} = \frac{8}{3}$.

2.2 The equation $x^2 + 6x + 4 = 0$ has roots α, β . Without solving it, form a quadratic equation with roots 3α and 3β . [3]

2.2 New sum $= 3(\alpha + \beta) = -18$; new product $= 9\alpha\beta = 36$. Equation: $x^2 + 18x + 36 = 0$.

2.3 For $x^2 - 5x + 9 = 0$ with roots α, β , find the value of $\frac{\alpha}{\beta} + \frac{\beta}{\alpha}$. [3]

2.3
$$\frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} = \frac{25 - 18}{9} = \frac{7}{9}.$$

Exercises 2.4·3.1

2.4 (*Original.*) $x^2 + px + q = 0$ has roots α and $\alpha + 2$. Show that $p^2 - 4q = 4$ for every such pair. [3]

3.1 (*Original.*) $g(x) = 2x^2 - 6x + m = 0$ has roots α, β with $\alpha^2 + \beta^2 = 11$. (a) Find m .
(b) Form a quadratic equation with integer coefficients whose roots are α^3 and β^3 . [6]

Exercises 2.4·3.1

2.4 (*Original.*) $x^2 + px + q = 0$ has roots α and $\alpha + 2$. Show that $p^2 - 4q = 4$ for every such pair. [3]

2.4 Sum $2\alpha + 2 = -p$ and product $\alpha(\alpha + 2) = q$. Then $p^2 = 4(\alpha + 1)^2 = 4\alpha^2 + 8\alpha + 4$ and $4q = 4\alpha^2 + 8\alpha$, so $p^2 - 4q = 4$ ✓ – independent of α .

3.1 (*Original.*) $g(x) = 2x^2 - 6x + m = 0$ has roots α, β with $\alpha^2 + \beta^2 = 11$. (a) Find m . (b) Form a quadratic equation with integer coefficients whose roots are α^3 and β^3 . [6]

3.1 (a) $(\alpha + \beta)^2 - 2\alpha\beta = 9 - m = 11 \Rightarrow m = -2$. (b) $\alpha\beta = -1$; $\alpha^3 + \beta^3 = 27 - 3(-1)(3) = 36$; $\alpha^3\beta^3 = (-1)^3 = -1$. Equation: $x^2 - 36x - 1 = 0$.

Exercise 3.2

3.2 (*Adapted from Edexcel IAL WFM01 June 2025 Q6.*) $f(x) = 3x^2 + kx - 5$ has roots α, β with $\alpha + \beta = 9\alpha\beta$. (a) State $\alpha\beta$ and find k . (b) Prove that $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ by expanding $(\alpha + \beta)^3$. (c) Without solving $f(x) = 0$, find a quadratic equation with integer coefficients whose roots are $\alpha^2 + \beta$ and $\alpha + \beta^2$. [10]

Exercise 3.2

3.2 (*Adapted from Edexcel IAL WFM01 June 2025 Q6.*) $f(x) = 3x^2 + kx - 5$ has roots α, β with $\alpha + \beta = 9\alpha\beta$. (a) State $\alpha\beta$ and find k . (b) Prove that $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ by expanding $(\alpha + \beta)^3$. (c) Without solving $f(x) = 0$, find a quadratic equation with integer coefficients whose roots are $\alpha^2 + \beta$ and $\alpha + \beta^2$. [10]

3.2 (a) $\alpha\beta = -\frac{5}{3}$; $-\frac{k}{3} = 9(-\frac{5}{3}) = -15 \Rightarrow k = 45$, $\alpha + \beta = -15$. (b) $(\alpha + \beta)^3 = \alpha^3 + 3\alpha^2\beta + 3\alpha\beta^2 + \beta^3 = (\alpha^3 + \beta^3) + 3\alpha\beta(\alpha + \beta)$; rearrange. (c) Sum = $\frac{640}{3}$, product = $-\frac{31040}{9}$ (see worked example): $9x^2 - 1920x - 31040 = 0$.

Local review guidance: symmetric identities applied without finding individual roots; exact fractions kept until the final integer equation; new equations presented as $x^2 - (\text{sum})x + \text{product} = 0$. This is authored classroom guidance.