

4.6 Vectors

Name: _____ Class: _____ Date: _____ **Total: 21 marks**

KEY WORDS magnitude • unit vector • position vector
• scalar product

Objective

Work with vectors in two and three dimensions: magnitude and unit vectors, addition and scalar multiplication with their geometric meanings, position vectors, and angles from the scalar product. Original teaching questions for specification items P4.7.1 to P4.7.7; the exam-style part adapts reviewed WMA14 June 2025 demand.

1 Key knowledge and worked examples

- Column and **ijk** forms: $a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$.
- Magnitude** : $|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$; **unit vector** $\hat{\mathbf{a}} = \mathbf{a}/|\mathbf{a}|$.
- Position vector** : $\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$.
- Scalar product** : $\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3 = |\mathbf{a}||\mathbf{b}|\cos\theta$. Parallel vectors are scalar multiples; perpendicular ones have zero scalar product.

Worked example – magnitudes and a unit vector. $\mathbf{a} = 2\mathbf{i} - \mathbf{j} + 2\mathbf{k}$.

- $|\mathbf{a}| = \sqrt{4 + 1 + 4} = 3$, so the unit vector is $\frac{1}{3}(2\mathbf{i} - \mathbf{j} + 2\mathbf{k})$.
- Check: its magnitude is $\frac{1}{3}\sqrt{9} = 1$ ✓.

Worked example – a 45° angle (Edexcel IAL WMA14 June 2025 Q6). $A = \mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$, $B = 5\mathbf{i} + 2\mathbf{j} + \mathbf{k}$, $C = 3\mathbf{i} + \alpha\mathbf{j} + 5\mathbf{k}$, angle $BAC = 45^\circ$; find α .

- $\overrightarrow{AB} = 4\mathbf{i} + 4\mathbf{k}$ with $|\overrightarrow{AB}| = \sqrt{32}$; $\overrightarrow{AC} = 2\mathbf{i} + (\alpha - 2)\mathbf{j} + 8\mathbf{k}$; scalar product $= 8 + 0 + 32 = 40$.

$$\cos 45^\circ = \frac{40}{\sqrt{32}\sqrt{68 + (\alpha - 2)^2}} \implies \sqrt{68 + (\alpha - 2)^2} = \frac{40\sqrt{2}}{4\sqrt{2}} = 10 \implies (\alpha - 2)^2 = 32$$

$$\alpha = 2 \pm 4\sqrt{2}$$

- Check: $|\overrightarrow{AB}||\overrightarrow{AC}|\cos 45^\circ = 4\sqrt{2} \times 10 \times \frac{1}{\sqrt{2}} = 40$ ✓.

2 Practice

2.1 Find $|3\mathbf{i} + 4\mathbf{j}|$ and the unit vector in that direction. [2]

2.2 For $P(1, 2, 3)$ and $Q(4, 6, 3)$, find $\overrightarrow{PQ}$ and its magnitude. [3]

2.3 Determine whether $\mathbf{a} = 2\mathbf{i} + (-1)\mathbf{j} + 3\mathbf{k}$ and $\mathbf{b} = (-4)\mathbf{i} + 2\mathbf{j} + (-6)\mathbf{k}$ are parallel. [2]

2.4 Find the angle between $1\mathbf{i} + 1\mathbf{k}$ and $1\mathbf{j} + 1\mathbf{k}$, giving an exact value. [4]

3 Exam-style questions

3.1 (Adapted from Edexcel IAL WMA14 June 2025 Q6.) Relative to O : A has position vector $\mathbf{i} - \mathbf{j} + 2\mathbf{k}$ and B has position vector $3\mathbf{i} + \mathbf{j} + 6\mathbf{k}$. The point C has position vector $2\mathbf{i} + \beta\mathbf{j} + 4\mathbf{k}$, and angle BAC is 60° . Find the exact possible values of β . [6]

3.2 The line segment RS has midpoint $M(1, 1, 0)$; R has position vector $3\mathbf{i} - \mathbf{j} + 4\mathbf{k}$. Find $\overrightarrow{RS}$ and the distance RS . [4]
