# 5.2 Complex roots of cubics and quartics

Name: _____ Class: _____ Date: _____ **Total: 28 marks**

**KEY WORDS** conjugate pair • quadratic factor • discriminant

## Complex roots of cubics and quartics

## Objective

Use the conjugate-root property to find all roots of cubic and quartic equations with real (integer) coefficients, building the real quadratic factor and dividing it out (FP1 1.5–1.6).

## 1 Key knowledge and worked examples

**Conjugate pairs.** If $f(z_1) = 0$ for a polynomial with real coefficients, then $f(\bar{z}_1) = 0$. From a root $p \pm qi$ build the factor $z^2 - 2pz + (p^2 + q^2)$ – sum $2p$, product $p^2 + q^2$ – then divide or compare coefficients to finish.

**Worked example – a quartic with one complex root (WFM01 June 2025 Q7).** $f(z) = Pz^4 - 36z^3 + Qz^2 + 192z + 68$ has root $3 + 5i$. The conjugate $3 - 5i$ is also a root, giving the factor $z^2 - 6z + 34$. Writing $f(z) = (z^2 - 6z + 34)(Pz^2 + bz + c)$: $34c = 68 \Rightarrow c = 2$; $-6c + 34b = 192 \Rightarrow b = 6$; $b - 6P = -36 \Rightarrow P = 7$; $c - 6b + 34P = Q \Rightarrow Q = 204$. The remaining quadratic $7z^2 + 6z + 2$ has discriminant $36 - 56 = -20$, so the other roots are $z = \dfrac{-3 \pm i\sqrt{5}}{7}$.

**Worked example – a cubic from one real root.** $g(x) = x^3 - 7x^2 + 17x - 15$ has $x = 3$ as a root. Divide: $g(x) = (x - 3)(x^2 - 4x + 5)$, and the quadratic has discriminant $16 - 20 < 0$, so the roots are $x = 3,\ 2 \pm i$. Check the product of roots: $3(2+i)(2-i) = 3 \cdot 5 = 15$ ✓ (constant $-15$ with leading coefficient 1 needs $\alpha\beta\gamma = 15$).

## 2 Practice

**2.1** Given that $2 + i$ is a root of $x^2 + px + q = 0$ with $p, q$ real, find $p$ and $q$. [2]

**2.2** $f(x) = x^3 - 3x^2 + 7x - 5$ has root $x = 1$. Find all roots of $f(x) = 0$. [4]

**2.3** One root of $x^4 - 2x^3 + 6x^2 - 2x + 5 = 0$ is $1 + 2i$. Find the other three roots. [5]

**2.4** A quartic with real coefficients has roots $1 \pm 3i$ and a repeated real root $k$, with constant term 50. Find $k$. [3]

## 3 Exam-style questions

**3.1** *(Original.)* $h(z) = 2z^4 + 2z^3 + 7z^2 + 4z + 15$ has a root $\frac{1}{2} + \frac{3}{2}i$. (a) Write down the real quadratic factor coming from this root. (b) Given $h(z) = (2z^2 - 2z + 5)(z^2 + mz + n)$, find $m$ and $n$. (c) Hence solve $h(z) = 0$ completely. [6]

**3.2** *(Adapted from Edexcel IAL WFM01 June 2025 Q7.)* $f(z) = Pz^4 - 36z^3 + Qz^2 + 192z + 68$ has root $3 + 5i$. Without using a calculator, determine $P$, $Q$ and the three other roots, giving them in simplest form. [8]