

Further Pure Mathematics 2

Pearson Edexcel · International A-Level · Pure Mathematics

Further Pure Mathematics 2: inequalities, further complex numbers and differential equations

Further Pure Mathematics 2 (FP2, paper WFM02/01, 1 hour 30 minutes, 75 marks, first assessment June 2020) builds directly on P1–P4 and FP1 and counts one sixth of an International Advanced Level. Every unit is equally weighted, so FP2 material can also surface inside Further Mathematics. P1–P4 assumed content stays live: partial fractions, parametric and implicit differentiation, integration by parts and substitution, and the whole FP1 complex-number and proof toolkit. This reference covers modulus inequalities, the method of differences, Euler’s relation and de Moivre’s theorem, loci and $z \rightarrow w$ transformations, first and second order differential equations, Maclaurin and Taylor series, and polar coordinates.

Must-know for FP2 (assumed, not re-taught):

Family	Formulae
modulus	$ x = x$ for $x \geq 0$ and $-x$ for $x < 0$; $ z = \sqrt{a^2 + b^2}$, $ z_1 z_2 = z_1 z_2 $, $\arg(z_1 z_2) = \arg z_1 + \arg z_2$
sums	$\sum_{r=1}^n r = \frac{1}{2}n(n+1)$, $\sum r^2$, $\sum r^3$ (booklet)
algebra	partial fractions, e.g. $\frac{1}{r(r+1)} = \frac{1}{r} - \frac{1}{r+1}$ (P4)
calculus	product, quotient and chain rules; integration by substitution, parts, partial fractions (P4)
differential equations	separation of variables; exponential growth/decay $y = Ae^{kt}$ (P4)

Inequalities involving the modulus sign


Two safe moves and one banned move. Safe: split into cases from the definition of the modulus ($|f| = f$ where $f \geq 0$, $-f$ where $f < 0$), and square both sides of $|A| > |B|$ to get $A^2 > B^2$. Banned: multiplying or dividing by an expression whose sign you have not fixed – case-split on that sign instead, and intersect each case with its own domain.

Worked example – one branch dies (specimen style). Solve $|x^2 - 1| > 2(x + 1)$. Where $x^2 \geq 1$: $x^2 - 2x - 3 > 0$, so $(x - 3)(x + 1) > 0$, giving $x > 3$ or $x < -1$; keeping only $x \leq -1$ or $x \geq 1$ leaves $x < -1$ or $x > 3$. Where $x^2 < 1$: $1 - x^2 > 2x + 2$ becomes $(x + 1)^2 < 0$ – impossible. The solution is $x < -1$ or $x > 3$. A dead branch is a result, not a mistake: report it.

Worked example – modulus on both sides (WFM02 June 2025 Q8). The curve C has equation $y = \frac{15x}{|x|+4}$ and the line l has equation $y = x - 2$.

(a) Solve $x - 2 > \frac{15x}{|x|+4}$. The denominator is always positive, so multiply straight through. For $x \geq 0$: $(x - 2)(x + 4) > 15x$ gives $x^2 - 13x - 8 > 0$, so $x > \frac{13+\sqrt{201}}{2}$ (the other root is negative). For $x < 0$: $(x - 2)(4 - x) > 15x$ gives $-x^2 - 9x - 8 > 0$, i.e. $(x + 1)(x + 8) < 0$, so $-8 < x < -1$. Together: $-8 < x < -1$ or $x > \frac{13+\sqrt{201}}{2}$.

(b) Solve $|x - 2| > \left| \frac{15x}{|x|+4} \right|$. Squaring (both sides non-negative) turns this into $(x - 2)(|x| + 4) > 15x$ – part (a) – or $(x - 2)(|x| + 4) < -15x$. The new branch: for $x \geq 0$, $x^2 + 17x - 8 < 0$ gives $0 \leq x < \frac{\sqrt{321}-17}{2}$; for $x < 0$, $x^2 - 21x + 8 > 0$ holds for every $x < 0$. So the answer is $x < \frac{\sqrt{321}-17}{2}$ or $x > \frac{13+\sqrt{201}}{2}$. The "hence" is real: half of part (b) is part (a)'s answer.



The method of differences

The **method of differences** sums a series by splitting the summand (usually via **partial fractions**) so that each term cancels part of a later one – the middle collapses and only the first and last fragments survive. Write out $r = 1$, $r = 2$, $r = n - 1$, $r = n$ explicitly and watch what cancels; that display earns the marks.

Worked example – telescoping to a closed form (WFM02 June 2025 Q5).

Show $\sum_{r=1}^n \frac{1}{(r+1)(r+2)} = \frac{n}{2(n+2)}$. Split:

$$\frac{1}{(r+1)(r+2)} = \frac{1}{r+1} - \frac{1}{r+2}.$$

The sum is $\left(\frac{1}{2} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \cdots + \left(\frac{1}{n+1} - \frac{1}{n+2}\right)$: everything from $\frac{1}{3}$ to $\frac{1}{n+1}$ cancels,

leaving $\frac{1}{2} - \frac{1}{n+2} = \frac{n}{2(n+2)}$. (Check $n = 1$: both sides $\frac{1}{6}$.) This is also how inequality follow-ups are set: $\frac{n}{2(n+2)} > 0.49$ needs $n > 98$, so the smallest is $n = 99$ – the algebra, not a table of values.

Euler's relation and de Moivre's theorem

Euler's relation $e^{i\theta} = \cos \theta + i \sin \theta$ turns the polar form into an exponential: $z = re^{i\theta}$, and $z^n = r^n e^{in\theta}$ becomes pure arithmetic. Read off the companions $\cos \theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta})$ and $\sin \theta = \frac{1}{2i}(e^{i\theta} - e^{-i\theta})$. **De Moivre's theorem**

$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$ follows, and is provable for every positive integer n by induction: true for $n = 1$; if true for k , multiplying $(\cos k\theta + i \sin k\theta)(\cos \theta + i \sin \theta)$ and expanding with the addition formulae gives $\cos(k+1)\theta + i \sin(k+1)\theta$. For negative n , take reciprocals and note $\cos(-n\theta) + i \sin(-n\theta)$ has modulus 1 and the right argument.

Worked example – a multiple-angle identity (WFM02 June 2025 Q9). Expand the imaginary part of $(\cos \theta + i \sin \theta)^5$ by the binomial theorem:

$$\text{Im} = 5c^4s - 10c^2s^3 + s^5 \quad (c = \cos \theta, s = \sin \theta).$$

De Moivre says this equals $\sin 5\theta$. Substituting $c^2 = 1 - s^2$: $5s(1 - s^2)^2 - 10s^3(1 - s^2) + s^5 = 16s^5 - 20s^3 + 5s$, so

$$\sin 5\theta = 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta.$$

Setting $5\theta = \pi$ gives $16s^4 - 20s^2 + 5 = 0$, so $s^2 = \frac{5 \pm \sqrt{5}}{8}$ and $\sin \frac{\pi}{5} = \sqrt{\frac{5 - \sqrt{5}}{8}} = \frac{\sqrt{10 - 2\sqrt{5}}}{4}$ – take the smaller root because $\sin \frac{\pi}{5} < \sin \frac{3\pi}{5}$, and indeed $\sin \frac{3\pi}{5}$ is the larger root.

Worked example – fifth roots (WFM02 June 2025 Q1). Solve $z^5 = -2\sqrt{2} - 2\sqrt{6}i$. Modulus $\sqrt{8 + 24} = \sqrt{32} = 4\sqrt{2}$; the argument is $-\frac{2\pi}{3}$ (third quadrant, $\tan = \sqrt{3}$). So

$$z = (4\sqrt{2})^{1/5} e^{i(-2\pi/15 + 2\pi k/5)} = \sqrt{2} e^{i(-\frac{2\pi}{15} + \frac{2\pi k}{5})}, \quad k = 0, 1, 2, 3, 4,$$

five roots spaced $\frac{2\pi}{5}$ apart on the circle $|z| = \sqrt{2}$. Use the n th-root pattern $\sqrt[n]{r} e^{i(\theta + 2\pi k)/n}$ – never decimal approximations.

Loci, regions and z→w transformations

Locus toolkit in the Argand diagram: $|z - a| = b$ is a circle (centre a , radius b); $|z - a| = |z - b|$ is the **perpendicular bisector** of a and b ; $|z - a| = k|z - b|$ with $k \neq 1$ is a circle (square both sides and collect terms); $\arg(z - a) = \beta$ is a half-line from a . Regions replace = by $<$ or $>$: test one convenient point (the origin rarely lies

on the locus). For **Möbius transformations**



$$w = \frac{az + b}{cz + d}, \text{ never try to}$$

substitute point-by-point: make z the subject first, then take moduli (or arguments) of both sides. A circle through the pole $z = -\frac{d}{c}$ maps to a straight line; a circle not through the pole maps to a circle.

Worked example – image of a circle (WFM02 June 2025 Q3). $w = \frac{9i - iz}{1 + z}$ ($z \neq -1$) maps P, where $|z| = 1$, to a line l . Making z the subject: $z(w + i) = 9i - w$, so $z = \frac{9i - w}{w + i}$. Then $|z| = 1$ forces

$$|9i - w| = |w + i| \implies u^2 + (v - 9)^2 = u^2 + (v + 1)^2 \implies 81 - 18v = 2v + 1,$$

so $20v = 80$ and l is $v = 4$ – the perpendicular bisector of $9i$ and $-i$, exactly as the equal-moduli reading predicts. Spot-check with the endpoints of the vertical diameter: $z = i \mapsto 5 + 4i$ and $z = -i \mapsto -5 + 4i$; both land on $v = 4$.



First order differential equations

Separable equations are assumed. The new standard form is $\frac{dy}{dx} + Py = Q$ with P, Q functions of x : multiply by the **integrating factor** $e^{\int P dx}$, chosen so the left side becomes $\frac{d}{dx} (ye^{\int P dx})$, then integrate once. For a **given substitution** (say $z = f(y)$), compute $\frac{dz}{dx}$ by the chain rule, rewrite the equation in z and x , solve the (now linear or separable) equation, and substitute back – the question names the target form, so end in it.

Worked example – integrating factor warm-up. $\frac{dy}{dx} + 2y = e^{-x}$: $e^{\int 2 dx} = e^{2x}$, so $\frac{d}{dx}(ye^{2x}) = e^x$ and $ye^{2x} = e^x + C$, i.e. $y = e^{-x} + Ce^{-2x}$.

Worked example – a substitution then an integrating factor (WFM02 June 2025 Q6). Equation (I) is $2\frac{dy}{dx} + y \cot x + y^3 \tan x \sec x = 0$ on $0 < x < \frac{\pi}{2}$.

(a) With $z = \frac{1}{y^2}$, $\frac{dz}{dx} = -2y^{-3}\frac{dy}{dx}$, so $2\frac{dy}{dx} = -y^3\frac{dz}{dx}$. Substituting and dividing by y^3 : $-\frac{dz}{dx} + y^{-2} \cot x + \tan x \sec x = 0$, i.e.

$$\frac{dz}{dx} - z \cot x = \tan x \sec x \quad (\text{II}).$$

(b) The integrating factor is $e^{-\int \cot x dx} = \csc x$, so $\frac{d}{dx}(z \csc x) = \csc x \tan x \sec x = \frac{1}{\cos^2 x} = \sec^2 x$. Hence $z \csc x = \tan x + C$, giving $z = \sin x \tan x + C \sin x$ and

$$y^2 = \frac{1}{\sin x \tan x + C \sin x}.$$

(c) At $x = \frac{\pi}{6}$: $y^2 = \frac{4\sqrt{3}}{3}$ so $z = \frac{\sqrt{3}}{4} = \frac{1}{2} \cdot \frac{1}{\sqrt{3}} + \frac{C}{2}$, forcing $C = \frac{1}{2\sqrt{3}}$. At $x = \frac{\pi}{3}$: $z = \frac{\sqrt{3}}{2} \cdot \sqrt{3} + \frac{1}{2\sqrt{3}} \cdot \frac{\sqrt{3}}{2} = \frac{3}{2} + \frac{1}{4} = \frac{7}{4}$, so $y^2 = \frac{4}{7}$ and $y = \pm \frac{2}{\sqrt{7}}$. Both signs are answers – y^2 , not y , was determined.

Second order differential equations

For $a\frac{d^2y}{dx^2} + b\frac{dy}{dx} + cy = f(x)$, the general solution is CF + PI: the **complementary function** solves the homogeneous equation via the **auxiliary equation** $am^2 + bm + c = 0$; the **particular integral** is any single solution of the full equation.

Auxiliary equation	Complementary function
distinct real roots $m_1 \neq m_2$	$y = Ae^{m_1x} + Be^{m_2x}$
repeated root m	$y = (A + Bx)e^{mx}$
complex roots $p \pm qi$	$y = e^{px}(A \cos qx + B \sin qx)$

Right-hand side $f(x)$	Trial PI
ke^{px}	λe^{px} ; if p is a root, use $\lambda x e^{px}$ (again if repeated)
$a + bx(+cx^2)$	$\lambda + \mu x(+\nu x^2)$ – every degree present
$m \cos \omega x + n \sin \omega x$	$\lambda \cos \omega x + \mu \sin \omega x$

Given initial or boundary conditions, find the constants last, from the full general solution.

Worked example – resonance, then conditions (WFM02 June 2025 Q4).

Given that $y = \lambda x e^{3x}$ is a particular integral of $4 \frac{d^2 y}{dx^2} - 11 \frac{dy}{dx} - 3y = 78e^{3x}$:

(a) $y' = \lambda e^{3x}(1 + 3x)$ and $y'' = \lambda e^{3x}(6 + 9x)$, so the left side is $\lambda e^{3x}(24 + 36x - 11 - 33x - 3x) = 13\lambda e^{3x}$. Matching $78e^{3x}$ gives $\lambda = 6$. (The plain trial λe^{3x} would give $0 \cdot e^{3x}$ – hence the extra x .)

(b) Auxiliary equation $4m^2 - 11m - 3 = (4m + 1)(m - 3) = 0$, so $m = -\frac{1}{4}$ or 3 : $y = Ae^{-x/4} + Be^{3x} + 6xe^{3x}$.

(c) With $y = \frac{9}{2}$ and $y' = 0$ at $x = 0$: $A + B = \frac{9}{2}$ and $-\frac{A}{4} + 3B + 6 = 0$. Solving, $A = 6$, $B = -\frac{3}{2}$, so

$$y = 6e^{-x/4} - \frac{3}{2}e^{3x} + 6xe^{3x}.$$

Maclaurin and Taylor series

The **Maclaurin series**

of f is $\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$ – differentiate as many

times as the question asks (third and higher derivatives are examinable in their own right). The **Taylor series**

about $x = a$ replaces x by the offset:

$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$. For a differential equation, differentiate the equation itself to climb the derivative ladder: $y(a)$ and $y'(a)$ feed the equation for $y''(a)$, then the derivative of the equation for $y'''(a)$, and so on – never re-derive, differentiate.

Worked example – a series solution about a non-zero centre (WFM02 June

2025 Q2). Given $\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} - 5y = e^{2x/9}$ with $y = 2$ and $y' = -1$ at $x = 3$:

at $x = 3$: $y'' = 4y' + 5y + e^{2x/9} = -4 + 10 + e^{2/3} = 6 + e^{2/3}$, so the quadratic term is $\frac{1}{2}(6 + e^{2/3})(x - 3)^2 = (3 + \frac{1}{2}e^{2/3})(x - 3)^2$. Differentiating the equation, $y''' = 4y'' + 5y' + \frac{2}{9}e^{2x/9}$, so at $x = 3$: $y''' = 4(6 + e^{2/3}) - 5 + \frac{2}{9}e^{2/3} = 19 + \frac{38}{9}e^{2/3}$, and the cubic term is $\frac{1}{6}y'''(x - 3)^3$. The series is

$$y = 2 - (x - 3) + \left(3 + \frac{1}{2}e^{2/3}\right)(x - 3)^2 + \left(\frac{19}{6} + \frac{19}{27}e^{2/3}\right)(x - 3)^3.$$

Polar coordinates and areas

In **polar coordinates**




a point is (r, θ) with $r \geq 0$ measured from the **initial**

line (the positive x -axis, θ anticlockwise). Convert with $x = r \cos \theta$, $y = r \sin \theta$,

$r^2 = x^2 + y^2$: for example $r = 4 \cos \theta$ gives $r^2 = 4r \cos \theta$, i.e. $x^2 + y^2 = 4x$ – the circle $(x - 2)^2 + y^2 = 4$. Know the sketch staples: $r = 2a \cos \theta$ (circle on the initial line

through the pole), $r = k\theta$ (spiral), $r = a(1 + \cos \theta)$ (**cardioid**), $r = a \cos 2\theta$ (four petals), $r^2 = a^2 \cos 2\theta$ (**lemniscate**), existing only where $\cos 2\theta \geq 0$). The area swept from $\theta = \alpha$ to β is $\frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$; tangents parallel to the initial line occur where $\frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta = 0$.

Worked example – exact areas. The whole **lemniscate** $r^2 = 8 \cos 2\theta$: by symmetry the right loop is $\frac{1}{2} \int_{-\pi/4}^{\pi/4} 8 \cos 2\theta d\theta = 4 \left[\frac{1}{2} \sin 2\theta \right]_{-\pi/4}^{\pi/4} = 4$, so both loops enclose total area 8. The **cardioid** $r = 2(1 + \cos \theta)$: $\frac{1}{2} \int_{-\pi}^{\pi} 4(1 + \cos \theta)^2 d\theta = 2 \int_{-\pi}^{\pi} (1 + 2 \cos \theta + \cos^2 \theta) d\theta = 2(2\pi + 0 + \pi) = 6\pi$, since $\int_{-\pi}^{\pi} \cos^2 \theta d\theta = \pi$.



Checklist before you call this unit done

- Solve modulus inequalities by case split and by squaring; handle $|x|$ inside denominators; report dead branches; reuse an earlier part when "hence" appears.
- Sum finite series by partial fractions and cancellation, writing the $r = 1, 2, n - 1, n$ lines; close the follow-up inequality by algebra.
- Use $e^{i\theta}$, the exponential forms of $\cos$ and $\sin$, and de Moivre (proved for integer n) to expand multiple angles and powers, and to find all n th roots in exact polar form.
- Sketch circles, bisectors, Apollonius circles and argument half-lines; test regions; make z the subject of a Möbius transformation and equate moduli or arguments.
- Solve $\frac{dy}{dx} + Py = Q$ with an integrating factor; execute given substitutions and answer in the form demanded, minding $\pm$ from y^2 .
- Solve second order equations via auxiliary equation and trial PI (including the multiply-by- x resonance), then fit initial conditions.
- Build Maclaurin and Taylor series, including series solutions of differential equations by differentiating the equation.
- Sketch polar staples, convert to and from Cartesian, and evaluate exact areas with $\frac{1}{2} \int r^2 d\theta$.