

Pure Mathematics 2

Edexcel IAL P2 (WMA12/01); compulsory for IAS/IAL Pure Mathematics and
Mathematics

1. Unit P2 at a glance

1. Unit P2 at a glance

WMA12/01: 1 hour 30 minutes, 75 marks. P1 knowledge is assumed and may be tested. Calculator and yellow formulae booklet as for P1; the P2 must-know list ($\log_a x + \log_a y \equiv \log_a(xy)$, $\tan \theta \equiv \sin \theta / \cos \theta$, $\sin^2 \theta + \cos^2 \theta \equiv 1$, $\text{area} = \int y dx$) is not in the booklet.

2. Proof: exhaustion and counterexample

2. Proof: exhaustion and counterexample

A proof proceeds from assumptions through logic to a conclusion. Exhaustion checks every case ($m = 3q + 1$ or $3q + 2$ covers all non-multiples of 3); one counterexample disproves a claim ($2^3 + 1 = 9$ is not prime).

3. Worked: $m^2 + 3m + 2$ divisible by 3

3. Worked: $m^2 + 3m + 2$ divisible by 3

$(m + 1)(m + 2)$ with $m = 3q + 1$ gives $3(3q + 2)(q + 1)$; with $m = 3q + 2$ gives $3(q + 1)(3q + 4)$. Both cases are multiples of 3, which exhausts every non-multiple of 3.

4. Factor and remainder theorems

4. Factor and remainder theorems

$(x - b)$ is a factor $\iff f(b) = 0$; dividing by $(ax - b)$ leaves remainder $f(b/a)$. After division, $f(x) = (ax - b)(\text{quotient}) + \text{remainder}$.

5. Worked: two conditions, two unknowns

5. Worked: two conditions, two unknowns

$f(x) = ax^3 + bx^2 + 5x - 3$ with $(x + 3)$ a factor and remainder 7 dividing by $(2x - 1)$:
 $f(-3) = 0$ gives $b = 3a + 2$; $f(\frac{1}{2}) = 7$ gives $a + 2b = 60$; so $a = 8$, $b = 26$.

6. Circle geometry

6. Circle geometry

$(x - a)^2 + (y - b)^2 = r^2$; complete the square on the expanded form. Diameter endpoints give the centre as midpoint and r^2 as a quarter of the diameter length squared. Tangent $\perp$ radius; centre-to-chord perpendicular bisects the chord.

7. Worked: circle from a diameter

7. Worked: circle from a diameter

Endpoints $(-2, 5)$ and $(4, 15)$: centre $(1, 10)$, $d^2 = 36 + 100 = 136$ so $r^2 = 34$:
 $(x - 1)^2 + (y - 10)^2 = 34$. Nearest point to the x-axis: $(1, 10 - \sqrt{34})$.

8. Arithmetic and geometric series

8. Arithmetic and geometric series

$a_n = a + (n - 1)d$, $S_n = \frac{n}{2}\{2a + (n - 1)d\}$; $a_n = ar^{n-1}$, $S_n = \frac{a(1-r^n)}{1-r}$, $S_\infty = \frac{a}{1-r}$ for $|r| < 1$. Both sum-formula proofs should be known; Σ notation is expected.

9. Worked: logs find the term number

9. Worked: logs find the term number

$10(0.8)^{k-1} < 0.0005 \implies (0.8)^{k-1} < 0.00005 \implies (k-1) \log 0.8 < \log 0.00005$, and dividing by the negative $\log 0.8$ flips the sign: $k-1 > 44.38$, so $k = 46$.

10. Binomial expansion

10. Binomial expansion

$(a + bx)^n = a^n + \binom{n}{1} a^{n-1}(bx) + \binom{n}{2} a^{n-2}(bx)^2 + \dots$ with $\binom{n}{r} = \frac{n!}{r!(n-r)!}$. Match coefficients term by term: in $(2 + kx)(1 - 3x)^5$, the x^2 coefficient $180 - 15k$.

11. Log laws and domains

11. Log laws and domains

$\log_a x + \log_a y \equiv \log_a(xy)$, $\log_a x - \log_a y \equiv \log_a \frac{x}{y}$, $k \log_a x \equiv \log_a(x^k)$. Logs need positive arguments: test every root in the original equation before accepting it.

12. Worked: exact $(\log_3 2, 12)$ 12. Worked: exact $(\log_3 2, 12)$

$$3^{2x+1} = 6 \times 3^x \implies 3u^2 = 6u \text{ with } u = 3^x \implies 3^x = 2 \implies x = \log_3 2, \text{ and} \\ y = 6 \times 2 = 12.$$

13. Identities into equations

13. Identities into equations

$\tan \theta \equiv \sin \theta / \cos \theta$ and $\sin^2 \theta + \cos^2 \theta \equiv 1$ turn mixed equations into one function — often a hidden quadratic. Factor rather than divide: dividing by $\cos \theta$ loses candidates.

$$14. \text{ Worked: } 3\tan(2x + \pi/5) = \sqrt{3}$$

$$14. \text{ Worked: } 3\tan(2x + \pi/5) = \sqrt{3}$$

$$\tan(2x + \frac{\pi}{5}) = \frac{1}{\sqrt{3}} = \tan \frac{\pi}{6}, \text{ so } 2x + \frac{\pi}{5} = \frac{\pi}{6} + k\pi \text{ and } x = \frac{29\pi}{60} \text{ or } \frac{59\pi}{60} \text{ in } 0 \leq x < \pi.$$

15. Stationary points

15. Stationary points

$y' = 0$ at stationary points; classify with y'' (positive = minimum, negative = maximum) or the sign change of y' . Increasing where $y' > 0$, decreasing where $y' < 0$. In modelling, eliminate the second variable with the constraint first.

16. Worked: minimum surface

16. Worked: minimum surface

Trough: $3x^2L = 12$ so $L = 4/x^2$ and $S = 6x^2 + 5xL = 6x^2 + \frac{20}{x}$. Then $S' = 12x - \frac{20}{x^2} = 0 \implies x^3 = \frac{5}{3}$, $S = 25.3 \text{ m}^2$, and $S'' = 12 + \frac{40}{x^3} > 0$.

17. Definite integrals as areas

17. Definite integrals as areas

$\int_a^b y dx = F(b) - F(a)$ is the area under the curve for $y \geq 0$. Between a curve and a line, sketch first and integrate (upper – lower) between the intersections.

18. Worked: area 13.5 between curve and chord

18. Worked: area 13.5 between curve and chord

$y = 3x + \frac{16}{x^2} - 8$ with $P(1, 11)$, $Q(4, 5)$: line $y = -2x + 13$ gives area 24; the curve gives $\left[\frac{3}{2}x^2 - \frac{16}{x} - 8x\right]_1^4 = 10.5$; region $= 24 - 10.5 = 13.5$.

19. The trapezium rule

19. The trapezium rule

$\int_a^b y \, dx \approx \frac{h}{2} [y_0 + y_n + 2(y_1 + \cdots + y_{n-1})]$. Convex curve (bends upwards, like x^2): chords sit above it — overestimate. Concave (like $\sqrt{x}$): chords below — underestimate. More strips improve the estimate; scaled integrals scale the answer ($\int \frac{f}{2} = \frac{1}{2} \int f$).

Check yourself

Check yourself

Exhaustion with complete cases; counterexamples; factor/remainder theorems with $(ax - b)$; circles from any form plus the three properties; both series families with proofs and logs; binomial coefficients; log laws with domain checks; identity-reduced trig equations in either unit; stationary points classified and modelled; areas by definite integral and by the trapezium rule with the error direction.