

6.2 Probability

Name: _____ Class: _____ Date: _____

KEY WORDS conditional probability

<!-- Session 2 native-skill companion -->

Objective

Practise probability within the labelled native lessons. Public original tasks; locally authored answer guidance.

1 Key knowledge and worked examples

■ **Probability trees and outcomes · Foundation**

A probability lies between 0 and 1. Exhaustive, mutually exclusive outcomes have probabilities summing to 1. Multiply successive branch probabilities and add separate routes to an outcome.

$$P(RR) = P(R_1)P(R_2 \mid R_1)$$

A bag contains 3 red and 2 blue counters. With replacement, $P(\text{two red})=3/5 \times 3/5=9/25=0.36$. Without replacement, the red-red branch is $3/5 \times 2/4=0.3$. Label each branch before multiplying.

Mutually exclusive means no overlap; independent means that knowing one event does not change the other's probability. Two disjoint events with positive probability are not independent.

Use a frequency table or a simple tree before calculating. Formal conditional probability formulae are outside this Foundation/Core support lesson.

■ **Probability, trees and conditional reasoning · Higher**


Multiply along a tree branch and add disjoint branches. With replacement, the composition stays fixed. Conditional probability is $P(A \text{ given } B)=P(A \cap B)/P(B)$, for $P(B)>0$.

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}, \quad P(B) > 0$$

Without replacement, $P(\text{two red})=3/5 \times 2/4=3/10$. $P(\text{one of each})=3/5 \times 2/4+2/5 \times 3/4=3/5$. If $P(A \cap B)=0.12$ and $P(B)=0.3$, $P(A \text{ given } B)=0.4$.

Mutually exclusive means no overlap; independent means that knowing one event does not change the other's probability. Two disjoint events with positive probability are not independent.

A two-way table makes the restricted denominator visible. Before using a product $P(A)P(B)$, justify independence from the context or the supplied information.



Original native-lesson illustration; labels belong to its worked example.

2 Practice

2.1 (Foundation) Find the probability of two red from 3 red and 2 blue without replacement.

2.2 (Foundation) Find the probability of blue on a single draw from 3 red and 2 blue.

2.3 (Foundation) Find the probability of one red and one blue in that bag.

2.4 (Higher) Find $P(A \text{ given } B)$ if $P(A \cap B)=0.12$ and $P(B)=0.3$.

3 Exam-style questions

3.1 Decide whether this claim is true. Justify with a concrete calculation or counterexample: Mutually exclusive events with positive probabilities must be independent.
