7.1 Radioactivity and half-life

Name: _____ Class: _____ Date: _____ **Total: 23 marks**

Objective

Build the skills to answer exam questions on **radioactivity** : atoms and isotopes, alpha, beta and gamma radiation, nuclear equations, background radiation, half-life, and the uses and dangers of radiation.

You must be able to:

- describe alpha, beta and gamma radiation and their penetration
- complete nuclear equations
- use half-life to find activity and time
- explain the uses and dangers of radioactive sources

1 Key knowledge and worked examples

■ Radioactivity

radiation	what it is	stopped by	ionising
alpha α	2 protons + 2 neutrons, $^{4}_{2}\mathrm{He}$	a sheet of paper, a few cm of air	strongly
beta β^-	a fast electron, $^{0}_{-1}\mathrm{e}$	a few mm of aluminium	moderately
gamma γ	an electromagnetic wave	many cm of lead (only reduced)	weakly

- **Isotopes** have the same number of protons but different numbers of neutrons. $^{A}_{Z}\mathrm{X}$: A is the nucleon number, Z the proton number.
- **Half-life** is the time for the activity (or number of undecayed nuclei) to halve.

Worked example 1: alpha decay. Radium-226 ($^{226}_{88}\mathrm{Ra}$) gives off an alpha particle. Find the new nucleus.

1. Known: $A = 226$; $Z = 88$; alpha = $^{4}_{2}\mathrm{He}$.
2. Rule: top and bottom numbers balance.
3. Numbers: $226 - 4 = 222$; $88 - 2 = 86$.
4. Result: $^{222}_{86}\mathbf{Rn}$ (radon).

Worked example 2: half-life. A source has activity 800 Bq and half-life 3 hours. Find its activity after 9 hours.

1. Known: 800 Bq; half-life 3 h; 9 h.
2. Rule: $9 \div 3 = 3$ half-lives; halve three times.
3. Numbers: $800 \to 400 \to 200 \to 100$.
4. Result: 100 **Bq**.

Answer shapes: in beta decay a neutron turns into a proton, so Z goes up by 1 and A stays the same.

2 Practice

2.1 Carbon-12 and carbon-14 are isotopes. State what is the same and what is different in their nuclei. [2]



2.2 Give **two** sources of background radiation. [2]



2.3 Complete the equation for the beta decay of carbon-14: $^{14}_{6}\mathrm{C} \to {}^{?}_{?}\mathrm{N} + {}^{0}_{-1}\mathrm{e}$. [2]




3 Exam-style questions

3.1 A source is placed in front of a Geiger–Müller tube. Paper does not reduce the count rate, but 5 mm of aluminium reduces it to background level. Which radiation does the source give out? Explain. [3]



3.2 A student measures the count rate from a radioactive source every minute. The background count is 20 counts per minute. The graph shows the corrected count rate.

(a) Use the graph to find the half-life of the source. [2]



(b) The measured count rate at the start was 660 counts per minute. Find the corrected count rate after 3 half-lives. [3]

(c) Explain why the count rate must be corrected for background radiation, and why the readings scatter about the line. [3]

(d) A hospital uses a gamma source to kill cancer cells and a tracer to study a patient's kidneys. Explain which radiation and half-life is best for a tracer, and how staff keep their dose low. [6]

exercises.lol exercises.com.cn