

5.2 Ideal gas molecules

Name: _____ Class: _____ Date: _____

Total: 21 marks


Objective

Cover ideal gas molecules (5.15–5.22): molecular motion and pressure, absolute zero and the Kelvin scale, temperature and kinetic energy, and the two gas relationships. Original IGCSE-level practice.

1 Key knowledge and worked examples

■ Molecules make pressure

force on the wall:
pressure



molecules in random motion collide with the walls — each collision exerts a force; pressure is the total force per unit area

■ Kelvin and the two laws

$K = ^\circ C + 273$; **absolute zero** $-273\ ^\circ C = 0\ K$. Kelvin temperature $\propto$ average molecular kinetic energy.

$$p_1 V_1 = p_2 V_2 \text{ (constant } T) \quad \frac{p_1}{T_1} = \frac{p_2}{T_2} \text{ (constant } V)$$

Worked check. 200 cm^3 of gas at 150 kPa and 300 K is warmed to 600 K in the same volume: $p_2 = 150 \times 600/300 = 300\text{ kPa}$; then released to double the volume at that temperature: $p = 300 \times 200/400 = 150\text{ kPa}$.

2 Practice

2.1 State the value of absolute zero in $^{\circ}\text{C}$. [1]

2.2 Convert 50°C to kelvin and 0 K to $^{\circ}\text{C}$. [2]

2.3 A gas at 100 kPa in 6 dm^3 is compressed to 2 dm^3 at constant temperature. Calculate the new pressure. [2]

2.4 A fixed volume of gas at 200 kPa and 300 K is warmed to 450 K . Calculate the new pressure. [2]

2.5 State what happens to gas molecules' average speed when temperature rises. [1]

3 Exam-style questions

3.1 Explain, in terms of molecules, why a sealed can's pressure rises when it is heated. [3]

3.2 A sealed cylinder of gas at 180 kPa and 300 K is cooled to 200 K at constant volume. Calculate the new pressure, and state one assumption. [3]

3.3 Explain why pumping a bicycle tyre gets harder as it fills, using particle ideas. [3]

3.4 A balloon holds 12 dm^3 of gas at 300 kPa at depth. It rises and expands to 40 dm^3 at the same temperature. Calculate the gas pressure at the surface, stating one assumption. [4]
