

## 2.8 Standardise values and interpret a stated normal model

Name: \_\_\_\_\_ Class: \_\_\_\_\_ Date: \_\_\_\_\_ **Total: 24 marks**

**KEY WORDS** normal distribution 正态分布 • z-score 标准分 • standard deviation 标准差

### Objective

Standardise values and interpret a stated normal model. Original GAC004-aligned practice, not an official assessment. Show the method as well as the result. All applied scenarios are invented classroom problems.


### 1 Key knowledge and worked examples

■ **Key knowledge: normal probabilities depend on the model and its parameters**

A **normal distribution** 正态分布 is a continuous symmetric bell-shaped model with mean  $\mu$  and positive **standard deviation** 标准差  $\sigma$ . Standardising gives a **z-score** 标准分,  $z = (x - \mu)/\sigma$ . Total area is 1; individual exact values have zero probability in this continuous model.

Use the supplied standard-normal cumulative values  $\Phi(z) = P(Z \leq z)$ :  $\Phi(0) = 0.5000$ ,  $\Phi(1) = 0.8413$ ,  $\Phi(1.5) = 0.9332$ ,  $\Phi(2) = 0.9772$ . Symmetry gives  $\Phi(-z) = 1 - \Phi(z)$ . Interval probability is upper cumulative area minus lower cumulative area. These probabilities require a stated normal assumption; a mean and standard deviation alone do not prove that real data are normal.

■ **Worked example: a threshold is an area beyond a standardised value**



Known: an invented measurement model has  $\mu = 50$  and  $\sigma = 10$ . For values above 60, standardise the threshold.

$$z = \frac{x - \mu}{\sigma} = \frac{60 - 50}{10} = 1$$

Use the complement of the supplied cumulative area.

$$P(X > 60) = 1 - \Phi(1) = 1 - 0.8413 = 0.1587$$

This is a model probability, not a guarantee that exactly 15.87% of a finite sample exceeds 60.

### 2 Practice

**2.1** A normal model has mean 100 and standard deviation 15. Find the z-score of 130.[2]

---

---

---

---

**2.2** Using the supplied cumulative values, find the probability that a standard-normal value is greater than 1. [2]

---

---

---

---

**2.3** Use symmetry and the supplied values to find the probability that a standard-normal value lies between negative 1 and 1. [3]

---

---

---

---

---

---

**2.4** A normal model has mean 70 and standard deviation 8. Find the raw values corresponding to z-scores negative 1 and 1.5. Give the probability above the latter value. [3]

### 3 Exam-style questions

**3.1** A stated normal model for completion time has mean 40 minutes and standard deviation 5 minutes. (a) Find the probability of completion between 35 and 50 minutes. (b) For 200 independently modelled tasks, find the expected number exceeding 50 minutes. Explain why that expectation is not a guaranteed observed count. [6]

**3.2** Two stated normal score models are A with mean 60 and standard deviation 10, and B with mean 75 and standard deviation 5. A learner scores 80 on A and 85 on B. (a) Find both z-scores and compare relative position. (b) Use the supplied table to find each upper-tail probability. (c) For 500 modelled scores on each scale, find each expected count above those thresholds. (d) Explain two limits of using these calculations to claim equal ability on different tests. [8]