

1.2 Plan and interpret a controlled investigation

Name: _____ Class: _____ Date: _____

Total: 24 marks

KEY WORDS independent variable 自变量 • dependent variable 因变量

• control variables 控制变量

Objective

Plan and interpret a controlled investigation. This is original GAC013-aligned practice, not an official assessment. All supplied case data and review records are fictional; explicitly cited historical context is factual. The case data are invented for learning; they are not findings about a real school, country or population. Use the supplied evidence and do not invent missing observations.

1 Key knowledge and worked examples

■ **Key knowledge:** define what changes, what is measured and what stays comparable

The **independent variable** 自变量 is the factor deliberately changed. The **dependent variable** 因变量 is the measured response. **Control variables** 控制变量 are conditions kept comparable so they do not explain the difference instead. A testable hypothesis predicts a response that observations could challenge. Repeat trials to study variation and report the actual results; a neat average does not remove every source of uncertainty.

■ **Source A: text or task brief**


Fictional teacher-approved room-temperature paper investigation: compare water retained by paper types A and B. Each fresh square is 10 cm by 10 cm. Record its dry mass, immerse it for 60 s in the same water conditions, let it drain for 10 s without squeezing, then record wet mass using the same checked balance. Handle spills as instructed; use no electrical device near wet materials. The teacher provides the prepared results for this written task.

Recorded retained-water masses from three fresh squares of each type: A, 3.0 g, 3.2 g, 2.8 g; B, 4.2 g, 4.0 g, 4.4 g. Retained mass is wet mass minus dry mass. No other brands, immersion times or area sizes were tested.

■ **Source B: response or review record**

Review record: one student proposes squeezing type A before weighing but not B. Another writes: “B absorbed exactly 1.200000 g more in every trial, so it is always best for every cleaning job. We can discard any result below our prediction.” The class has not tested strength, cost, cleaning performance or other paper batches.

■ **Worked example: calculate a mean while keeping variation visible**



Known: type A retained 3.0 g, 3.2 g and 2.8 g. Why use a mean? It summarises these three trials without pretending they are identical.

$$\bar{m} = \frac{m_1 + m_2 + m_3}{3}$$

$$\bar{m}_A = \frac{3.0 + 3.2 + 2.8}{3} = 3.0 \text{ g}$$

The spread remains relevant. Keep the same draining method for both types; squeezing one would change the comparison.

2 Practice

2.1 Identify the independent variable and dependent variable.

[2]

2.2 State TWO conditions that must be kept comparable.

[2]

2.3 Calculate the mean for B and the difference between the two means, with units. [3]

2.4 Explain why squeezing A alone undermines the comparison and why repeats do not repair it. [3]

3 Exam-style questions

3.1 Evaluate the student’s result claim and data handling using six points. [6]

3.2 Write an 80–110-word investigation conclusion with a defensible follow-up. [8]
