

12.1 Algebra: domains, systems, inequalities and functions

Name: _____ Class: _____ Date: _____

KEY WORDS domain restriction • simultaneous solution

Objective

Solve equations while preserving denominator and radical restrictions; Analyse inequalities and simultaneous conditions without losing endpoints; Evaluate functions and translate verbal relationships into algebra.

Original public tasks and locally authored marking guidance. No official score is assigned.

1 Key knowledge and worked examples

domain restriction : A condition identifying inputs on which the original expression is defined.

simultaneous solution : Values satisfying every relation in a system at the same time.


Write restrictions before manipulating an equation. A denominator must be nonzero; a real even root needs a nonnegative argument. Cancelling factors preserves the expression only on its original domain. Squaring can introduce candidates because a square loses sign information; verify every candidate in the unsquared relation.

Solving a system requires the same ordered pair to satisfy all relations. Substitution or elimination can expose one, no or infinitely many linear solutions. For an inequality, multiplication or division by a negative reverses direction. For a factored quadratic inequality, order roots and test interval signs, including endpoints only when equality is allowed.

A function assigns one output to each permitted input. Evaluate the inner function first in a composition, preserving domain restrictions. Read a slope as change in output per input unit and an intercept as the output at input zero only when that input is meaningful. A verbal model should name variables and keep quantities in compatible units.

For Quantitative Comparison, preserve all allowed values. A variable with $x^2=4$ could be -2 or 2 unless constrained. One counterexample can reject a universal relationship; several favourable substitutions cannot prove it for every real value. Algebraic structure can establish a relationship without exhaustive numerical sampling.

For $\sqrt{x+2}=x$, $x \geq 0$. Squaring gives $x^2 - x - 2 = 0$, candidates 2 and -1 . Only 2 satisfies the original. For $(x-1)(x-4) \leq 0$, an upward quadratic is nonpositive on $1 \leq x \leq 4$, including roots. In the system $x+y=10$, $x-y=2$, adding gives $2x=12$; $x=6$ and $y=4$. For $f(t)=2t+1$ and $g(x)=x^2$, $f(g(3))=2 \cdot 9+1=19$, applying g first.



The value of a decides between one solution and infinitely many

2 Practice

2.1 Solve $(x^2 - 4)/(x - 2) = 5$ on its original real domain. Solve $(x + 1)(x - 3) \leq 0$ and find $f(g(2))$ for $f(t) = t^2$, $g(x) = x - 3$.

3 Exam-style questions

3.1 For real a , classify the system $x + y = 4$, $2x + ay = 8$ as having one, none or infinitely many solutions. Compare with the second equation $2x + ay = 9$. Give solutions in every consistent case.
