

# 31.1 Metric completeness and closure from a basis

Name: \_\_\_\_\_ Class: \_\_\_\_\_ Date: \_\_\_\_\_ **Total: 10 marks**

**KEY WORDS** complete metric space • neighbourhood basis

## Objective

Check a pullback metric using the properties of its defining map; Decide completeness through the image of an isometry; Determine closure using every basic neighbourhood instead of Euclidean intuition.

Original public practice. Marking guidance is authored locally. Use the topic reference for prerequisites and full theorem statements.


## 1 Key knowledge and worked examples

**complete metric space** : A metric space in which every Cauchy sequence converges to a point of that space.

**neighbourhood basis** : Basic neighbourhoods sufficient to test local topological properties.

For an injective map  $h:X\rightarrow R$ ,  $d(x,y)=|h(x)-h(y)|$  is a metric: symmetry and the triangle inequality come from  $R$ , and injectivity ensures  $d(x,y)=0$  only when  $x=y$ . The map  $h$  is an isometry onto its image  $h(X)$ . If  $h$  is not injective, this construction may be only a pseudometric. Bounded distances do not prove a space complete, and a familiar set of points can have very different Cauchy behaviour under different metrics.

Under  $d(x,y)=|\arctan x-\arctan y|$ , the integer sequence goes toward a missing image endpoint rather than a point of  $R$ , so it is Cauchy without convergence in that space. In the divisor basis, 16 lies in closure of  $\{8\}$ , since any basic set containing 16 also contains 8. But 4 does not:  $U_4=\{2,4\}$  is a neighbourhood of 4 missing 8. This explicitly separates multiples from divisors.



*The image of a Cauchy sequence can converge to a point outside the image*

## 2 Practice

**2.1** Which statement holds for  $d(x,y)=|\arctan x-\arctan y|$  on  $R$ ? [1]

- **A** It is not a metric
- **B** It is complete because  $\arctan$  is continuous
- **C** It is incomplete because its isometric image omits its endpoints
- **D** Every unbounded sequence fails to be Cauchy
- **E** Its distance equals ordinary absolute difference

**2.2** For the divisor basis in the lesson, which point belongs to closure of  $\{8\}$ ? [1]

- **A** 2
- **B** 3
- **C** 4
- **D** 6
- **E** 16

**2.3** On  $\mathbb{R}$  compare  $d_1(x,y)=|x^3-y^3|$  and  $d_2(x,y)=|\arctan x-\arctan y|$ . Prove they are metrics and decide completeness. [4]

3 Exam-style questions

3.1 On  $X = \{2, 3, 4, \dots\}$  use the divisor basis  $U_k = \{n \in X : n \mid k\}$ . Find the closure of  $\{6\}$  and the interior of  $\{6\}$ . Verify the basis conditions, including empty intersections. [4]

---

---

---

---

---

---

---

---

---

---

## Solutions

---

**2.1 C.** The injective map defines a metric. The image is an open bounded interval, and  $\arctan n$  tends to a missing endpoint. Injectivity proves this is a metric. Continuity does not give completeness, unbounded  $n$  is Cauchy in this metric, and arctangent distance is not ordinary absolute distance.

**2.2 E.** Closure consists of multiples of 8. The point 16 has every neighbourhood containing 8;  $U_4$  demonstrates why the divisor 4 does not belong. Closure requires multiples of eight. Two, three, four and six have divisor neighbourhoods missing eight; sixteen is a multiple and every neighbourhood contains eight.

**2.3** Both defining maps are injective, giving positive separation; absolute distance gives symmetry and the triangle inequality. Each map is an isometry onto its real image. The cube map has image all of  $\mathbb{R}$ , a complete space, so  $d_1$  is complete. The arctangent image is the open interval  $(-\pi/2, \pi/2)$ , which is incomplete. The sequence  $n$  is  $d_2$ -Cauchy because  $\arctan n$  tends to  $\pi/2$ , but no real point has that image, so it has no  $d_2$ -limit. Bounded metric values do not imply completeness.

**3.1** Every  $x$  belongs to  $U_x$ . An intersection is  $U_{\gcd(k,l)}$  if the gcd is at least two, and is empty for gcd one. Thus a point in an intersection has a basic neighbourhood within it. The smallest basic neighbourhood of  $x$  is  $U_x$ , contained in every  $U_k$  that contains  $x$ . It meets  $\{6\}$  exactly when  $6 \mid x$ . Therefore the closure is  $\{6, 12, 18, \dots\}$ , the multiples of six. The point six is not interior to its singleton because  $U_6$  also contains two and three. No other point is in that singleton, so its interior is empty. This closure calculation uses all basic neighbourhoods, not Euclidean intuition.

Local four-mark guidance for each written task: choose and justify the method; carry through the reasoning; give every requested result; check the relevant restriction, hypothesis or boundary case. These are classroom marks, not GRE scoring.