

4. A.2 · Abstract algebra and number theory

Starter Quiz ·

GRE Subject Test · GRE Mathematics

Name: _____ Score: _____

1. Which is a group?

- ☐ the integers under multiplication
- ☐ the nonzero rationals under multiplication
- ☐ the integers mod 8 without 0, under multiplication mod 8
- ☐ the positive integers under addition

2. Which residue has a multiplicative inverse modulo 9?

- ☐ 3
- ☐ 12
- ☐ 6
- ☐ 4

3. Find the least non-negative x with $5x \equiv 3 \pmod{7}$.

4. $\mathbb{Z}_7$ is an integral domain.

- ☐ True ☐ False

5. A homomorphism is injective exactly when its _____ is trivial.

6. What is the order of the element 4 in the additive group $\mathbb{Z}_{12}$?

7. For reduction $\varphi : \mathbb{Z}_{12} \rightarrow \mathbb{Z}_4$, how many elements are in the kernel?

8. Which ring is an integral domain under the convention that 1 and 0 are distinct?

- ☐ The zero ring
- ☐ $\mathbb{Z}_4$
- ☐ $\mathbb{Q}[x]$
- ☐ $\mathbb{Z}_6$

9. Find the least nonnegative inverse of 17 modulo 43.

10. How many distinct solution classes modulo 15 satisfy $6x \equiv 9 \pmod{15}$?

GRE Subject Test · GRE Mathematics

1. **the nonzero rationals under multiplication**

The others lack inverses (or closure).

2. **4**

$$\gcd(4, 9) = 1; 4 \cdot 7 = 28 \equiv 1.$$

3. **2**

$$5 \cdot 2 = 10 \equiv 3.$$

4. **True**

For $n \geq 2$, the ring $\mathbb{Z}_n$ has no nonzero zero divisors exactly when n is prime. Since 7 is prime, $\mathbb{Z}_7$ is a field and hence an integral domain.

5. **kernel**

Only the identity maps to the identity.

6. **3**

The successive multiples are 4, 8, 0 modulo 12. The least positive number of additions returning to 0 is 3.

7. **3**

The inputs sent to 0 modulo 4 are 0, 4 and 8. These are three distinct residues modulo 12.

8. **$\mathbb{Q}[x]$**

In $\mathbb{Q}[x]$, the product of nonzero polynomials has nonzero leading coefficient, so there are no zero divisors. In $\mathbb{Z}_6$, $2 \cdot 3 = 0$; in $\mathbb{Z}_4$, $2 \cdot 2 = 0$. The zero ring has $1 = 0$.

9. **38**

Bézout gives $1 = 2 \cdot 43 - 5 \cdot 17$. Thus -5 , or 38 modulo 43, is the inverse. Direct check: $17 \cdot 38 = 646 = 15 \cdot 43 + 1$.

10. **3**

The gcd is 3, which divides 9. Dividing both sides and the modulus gives $2x \equiv 3 \pmod{5}$, so $x \equiv 4 \pmod{5}$. The three classes modulo 15 are 4, 9 and 14.