

18. C.7 · Coordinate changes and vector integral theorems

Starter Quiz ·

GRE Subject Test · GRE Mathematics

Name: _____ Score: _____

1. The outward flux of $\mathbf{F} = (x, y, z)$ through the unit sphere is $k\pi$. Find k .
- _____
2. The counter-clockwise circulation of $(-y, x)$ around the circle of radius 3 is $k\pi$. Find k .
- _____
3. The spherical volume element (φ measured from the z -axis) is
- ☐ $\rho\,d\rho\,d\varphi\,d\theta$
- ☐ $\rho^2\sin\varphi\,d\rho\,d\varphi\,d\theta$
- ☐ $\sin\varphi\,d\rho\,d\varphi\,d\theta$
- ☐ $\rho^2\cos\varphi\,d\rho\,d\varphi\,d\theta$
4. A curl-free field on $\mathbb{R}^2 \setminus \{0\}$ is conservative.
- ☐ True ☐ False
5. $\mathbf{F} = \nabla(x^2y)$. Find $\int_C \mathbf{F} \cdot d\mathbf{r}$ along any path from $(0, 0)$ to $(2, 3)$.
- _____
6. The integral of $x^2 + y^2$ over $1 \leq x^2 + y^2 \leq 4$ is $k\pi$. Find k .
- _____
7. The volume of $0 \leq z \leq 4 - x^2 - y^2$ is $k\pi$. Find k .
- _____
8. If a field is conservative on an open connected domain, that domain must be simply connected.
- ☐ True ☐ False
9. The integral of $x^2 + y^2 + z^2$ over the ball of radius two is $k\pi$. Find k .
- _____
10. For $\mathbf{F} = (-y/2, x/2, z)$, the circulation around $x^2 + y^2 = 4, z = 0$, clockwise from above, is $k\pi$. Find k .
- _____

18. C.7 · Coordinate changes and vector integral theorems

Answers ·

GRE Subject Test · GRE Mathematics

1. **4**
- By divergence, outward flux is divergence 3 times unit-ball volume $4\pi/3$. Thus flux is 4π and $k = 4$.
2. **18**
- Green gives scalar curl $Q_x - P_y = 1 - (-1) = 2$. The radius-three disk has area 9π , so circulation is 18π and $k = 18$.
3. $\rho^2 \sin \varphi\,d\rho\,d\varphi\,d\theta$
- The Jacobian is $\rho^2 \sin \varphi$.
4. **False**
- The angular field circulates 2π .
5. **12**
- $\varphi(2, 3) - \varphi(0, 0) = 12$.
6. **7.5**
- The integrand is r^2 and the area factor is r . Thus $2\pi \int_1^2 r^3\,dr = 15\pi/2$, so $k = 7.5$.
7. **8**
- The projection disk has $0 \leq r \leq 2$. Integrating the height with the cylindrical factor gives $2\pi \int_0^2 (4-r^2)r\,dr = 8\pi$.
8. **False**
- On the punctured plane, $(2x, 2y)$ has the global potential $x^2 + y^2$. Simple connectedness is sufficient for the general zero-curl implication, not necessary for an individual field to be conservative.
9. **25.6**
- The integrand becomes ρ^2 and dV contributes another $\rho^2 \sin \varphi$. Angular integration gives 4π , and $\int_0^2 \rho^4\,d\rho = 32/5$, so $k = 128/5 = 25.6$.
10. **-4**
- The upward/counter-clockwise orientation gives 4π . Reversing traversal gives -4π , matching a downward surface normal.