

Starter Quiz

1. For every function f and sets A, B in its domain, which always holds?
- ☐ $f(A \cap B) = f(A) \cap f(B)$
- ☐ $f(A \setminus B) = f(A) \setminus f(B)$
- ☐ $f(A) \cap f(B) \subseteq f(A \cap B)$
- ☐ $f(A \cup B) = f(A) \cup f(B)$
2. How many equivalence classes does “same remainder mod 6” have on $\mathbb{Z}$?
- _____
3. On $\mathbb{Z}$, the relation “ $x - y$ is odd” is
- ☐ reflexive but not symmetric
- ☐ transitive and reflexive
- ☐ an equivalence relation
- ☐ symmetric but not reflexive
4. The negation of “for every $\varepsilon > 0$ there is N with $|a_n| < \varepsilon$ for all $n > N$ ” is
- ☐ For every $\varepsilon > 0$ and every cutoff N , some $n > N$ has $|a_n| \geq \varepsilon$.
- ☐ There is $\varepsilon > 0$ such that for every cutoff N , some $n > N$ has $|a_n| \geq \varepsilon$.
- ☐ There are $\varepsilon > 0$ and a cutoff N such that every $n > N$ has $|a_n| \geq \varepsilon$.
- ☐ For every $\varepsilon > 0$, some cutoff N makes every $n > N$ satisfy $|a_n| \geq \varepsilon$.
5. $f^{-1}(C \cap D) = f^{-1}(C) \cap f^{-1}(D)$ for every function f .
- ☐ True ☐ False
6. $P \Rightarrow Q$ is false only when P is true and Q is _____.
- _____
7. Let a, b be distinct and $f(a) = f(b) = L$. For $A = \{a\}$ and $B = \{b\}$, what are $f(A \cap B)$ and $f(A) \cap f(B)$?
- ☐ $\{L\}$ and $\emptyset$
- ☐ Both are $\emptyset$.
- ☐ $\emptyset$ and $\{L\}$
- ☐ Both are $\{L\}$.

9. Let $X = \{a, b, c\}$, $Y = \{L, M, N\}$, $f(a) = f(b) = L$, $f(c) = M$, and $C = \{L, N\}$. Find $f(f^{-1}(C))$.
- ☐ $\{a, b\}$
- ☐ $\{L, M\}$
- ☐ $\{L, N\}$
- ☐ $\{L\}$
10. The relation $xy > 0$ is an equivalence relation on all of $\mathbb{R}$.
- ☐ True ☐ False

1. $f(A \cup B) = f(A) \cup f(B)$

Union equality holds for every map. The other equalities/inclusion become valid for every pair under injectivity, but can fail for a map with a collision. A particular pair can still give equality without injectivity.

2. **6**

Two integers are related when they leave the same remainder on division by 6. The possible remainders 0, 1, ..., 5 give exactly 6 classes.

3. **symmetric but not reflexive**

It fails reflexivity because $x - x = 0$ is even. It is symmetric ($x - y$ odd $\Rightarrow y - x$ odd) but not transitive: $1 - 2$ and $2 - 3$ are odd, yet $1 - 3$ is even.

4. **There is $\varepsilon > 0$ such that for every cutoff N , some $n > N$ has $|a_n| \geq \varepsilon$.**

Flip $\forall \varepsilon, \exists N, \forall n > N$ to $\exists \varepsilon, \forall N, \exists n > N$, in that order. Failure of a strict inequality includes equality. Alternating zeros and ones fails at $\varepsilon = 1/2$ but has late zeros, and $\varepsilon = 2$ never fails.

5. **True**

An input maps into $C \cap D$ exactly when it maps into both C and D ; injectivity is unnecessary.

6. **false**

Truth table of implication.

7. $\emptyset$ and $\{L\}$

The input intersection is empty, but both image sets contain L .

8. **2**

$f(A) = \{4\}$, whose preimage is $\{-2, 2\}$. The round trip can add inputs.

9. $\{L\}$

Only L is in both C and the image $\{L, M\}$. No input maps to N .

10. **False**

At $x = 0$, x^2 is not positive, so reflexivity fails. Removing zero gives an equivalence relation with two sign classes.