

11.1 Polynomials and field extensions

Name: _____ Class: _____ Date: _____

Total: 10 marks**KEY WORDS** irreducible polynomial • extension degree

Objective

Test polynomial irreducibility over the specified field; Construct small finite fields from irreducible polynomials; Use extension degrees and the tower law; Use cyclotomic roots and coefficient relations to compute sums and products.

Original public practice. Marking guidance is authored locally. Use the topic reference for prerequisites and full theorem statements.

1 Key knowledge and worked examples

irreducible polynomial : A positive-degree polynomial with no factorisation into smaller positive degrees over the stated field.

extension degree : The dimension of an extension field as a vector space over its base field.

A polynomial is irreducible over a field if it has positive degree and no factorisation into polynomials of smaller positive degrees there. A quadratic or cubic is irreducible exactly when it has no root in that field. This root test alone fails for degree four or higher; for instance $(x^2+1)(x^2+2)$ has no real root but is reducible over $\mathbb{R}$. Always name the base field.

In $\mathbb{F}_2[\alpha]$ with $\alpha^2+\alpha+1=0$, multiply $\alpha(\alpha+1) = \alpha^2 + \alpha = (\alpha+1) + \alpha = 1$. Thus $\alpha^{-1}=\alpha+1$. Also $\alpha^3=1$ and α is not 1, so its multiplicative order is 3. This produces a field of four elements. By contrast $2 \cdot 2=0$ in $\mathbb{Z}/4\mathbb{Z}$, proving that quotient is not a field.

$F_2[\alpha]$, with $\alpha^2 = \alpha + 1$: all nonzero elements invertible

$\times$	1	α	$\alpha + 1$
1	1	α	$\alpha + 1$
α	α	$\alpha + 1$	1
$\alpha + 1$	$\alpha + 1$	1	α

Original course illustration; its values belong to the worked example, not the later practice.

2 Practice

2.1 Which polynomial is irreducible over F_2 ?

[1]

- A x^2
- B x^2+1
- C x^2+x
- D x^2+x+1
- E x^3+x^2

2.2 If $[L:K]=3$ and $[M:L]=2$, what is $[M:K]$?

[1]

- A 1
- B 2
- C 3
- D 5
- E 6

2.3 Construct $\mathbb{F}_2[x]/(x^2 + x + 1)$. List its elements and find the inverse of the residue $\alpha = [x]$. [4]

3 Exam-style questions

3.1 Is $x^4 + 4$ irreducible over $\mathbb{Q}$? Does the absence of rational roots answer that question? Can a degree-two extension of $\mathbb{Q}$ contain a root of an irreducible cubic over $\mathbb{Q}$? [4]

Solutions

2.1 D. The quadratic x^2+x+1 has no root among 0 and 1. The others factor as $x \cdot x$, $(x+1)^2$, $x(x+1)$ and $x^2(x+1)$, respectively, over F_2 . The four rejected polynomials factor respectively as x times x , $(x+1)$ squared, $x(x+1)$, and x squared times $(x+1)$ in characteristic two.

2.2 E. The tower law multiplies finite degrees, giving 2 times 3=6. One, two and three omit a nontrivial extension factor; five adds degrees instead of multiplying them.

2.3 The quadratic takes value one at both elements of $\mathbb{F}_2$, so it has no root and is irreducible. Its quotient is a field. Division leaves a representative of degree below two: $0, 1, \alpha, 1 + \alpha$. The relation is $\alpha^2 = \alpha + 1$ in characteristic two. Consequently $\alpha(\alpha + 1) = \alpha^2 + \alpha = 1$, so $\alpha^{-1} = \alpha + 1$. This is not $\mathbb{Z}/4\mathbb{Z}$, which has a nonzero zero divisor.

3.1 Direct multiplication gives $x^4 + 4 = (x^2 - 2x + 2)(x^2 + 2x + 2)$, so it is reducible over $\mathbb{Q}$ despite having no real, hence no rational, roots. The no-root test characterises irreducibility only for degree two or three over a field. If an irreducible cubic root β belonged to a degree-two extension L , then $\mathbb{Q} \subset \mathbb{Q}(\beta) \subset L$ and the tower law would make 3 divide 2. This is impossible. The conclusion uses irreducibility, not merely a cubic equation.

Local four-mark guidance for each written task: choose and justify the method; carry through the reasoning; give every requested result; check the relevant restriction, hypothesis or boundary case. These are classroom marks, not GRE scoring.