

21.1 Conditioning, Bayesian inference and sampling error

Name: _____ Class: _____ Date: _____

Total: 10 marks

KEY WORDS posterior probability • standard error

Objective

Separate conditional, joint and independent-event probabilities; Calculate posterior probabilities with a complete base-rate table; Use variance and sample size to determine a sample mean standard error.

Original public practice. Marking guidance is authored locally. Use the topic reference for prerequisites and full theorem statements.

1 Key knowledge and worked examples

posterior probability : A probability conditional on the observed evidence after accounting for prior proportions.

standard error : The standard deviation of a statistic across repeated samples.

For $P(B) > 0$, $P(A|B) = P(A \cap B) / P(B)$: restrict the population to B before computing the fraction in A. Independence means $P(A \cap B) = P(A)P(B)$, equivalently $P(A|B) = P(A)$ when the denominator is nonzero. Mutually exclusive events with positive probabilities cannot be independent, because their joint probability is zero. Sampling without replacement usually changes later probabilities; a fixed denominator for successive draws would describe a different experiment.

Let $P(A) = 0.4$, $P(B) = 0.5$ and $P(A \cap B) = 0.3$. Then $P(A|B) = 0.6$, so A and B are not independent, since $0.3 \neq 0.4 \cdot 0.5$. If a variable has standard deviation 12, an independent sample of size 36 has mean standard error 2. Increasing the sample size to 144 gives 1; the individual-observation standard deviation stays 12.

Condition on positives: 80 / 125, not 80 / 100

Expected group	Positive	Negative	Total
Defective	80	20	100
Not defective	45	855	900
Total	125	875	1000

Original course illustration; its values belong to the worked example, not the later practice.

2 Practice

2.1 $P(A)=0.4$, $P(B)=0.5$ and $P(A \cap B)=0.3$. What is $P(A|B)$?

[1]

- A 0.2
 - B 0.3
 - C 0.4
 - D 0.5
 - E 0.6
-

2.2 An iid sample of size 25 has mean standard error 6. What is the mean standard error for size 100 from the same population?

[1]

- A 1.5
 - B 3
 - C 6
 - D 12
 - E 24
-

2.3 In a stated model, 10% of devices have a defect. A test is positive for 80% of defective and 5% of nondefective devices. Find the probability of a defect given a positive result using 1000 expected devices. [4]

3 Exam-style questions

3.1 Independent observations have variance 25. Find the standard error of their mean for $n=100$. If instead every distinct pair has covariance 1, find the mean's variance. Does quadrupling n necessarily halve the standard error in that second model? [4]
