

1.1 Classical mechanics

Name: _____ Class: _____ Date: _____

Total: 12 marks

KEY WORDS Lagrangian • angular momentum

Objective

Apply Newtonian, Lagrangian and Hamiltonian descriptions; Analyse rotation, central forces and oscillations; Use conservation laws with their conditions. Original public practice; not an ETS paper.

1 Key knowledge and worked examples

Newton's second law is a vector statement. Choose coordinates and identify constraints before components.

Energy conservation applies when work from nonconservative forces is accounted for; momentum conservation needs zero net external impulse.

For small oscillations, linearise around a stable equilibrium. A restoring term proportional to displacement gives harmonic motion.

Lagrange's equations use $L=T-V$ and generalised coordinates. This undergraduate formalism is distinct from school-level force substitution.

■ Worked method

Choose an inertial frame and a coordinate x measured from spring equilibrium. For a conservative spring, the Lagrangian avoids solving constraint forces.

$$L = \frac{1}{2}m\dot{x}^2 - \frac{1}{2}kx^2.$$


$$\frac{d}{dt}\frac{\partial L}{\partial \dot{x}} - \frac{\partial L}{\partial x} = 0, \quad m\ddot{x} + kx = 0.$$

With $m = 2.0$ kg and $k = 18$ N/m,

$$\omega = \sqrt{k/m} = \sqrt{(18 \text{ N/m})/(2.0 \text{ kg})} = 3.0 \text{ rad/s}.$$

$$T = 2\pi/\omega = 2\pi/(3.0 \text{ rad/s}) = 2.09 \text{ s}.$$

Positive stiffness gives stable motion. The units of k/m are inverse seconds squared.



2 Practice

2.1 Choose one answer. Doubling spring mass changes angular frequency by: [1]

- A 2
- B $1/\sqrt{2}$
- C $1/2$

2.2 Choose one answer. In an isolated perfectly inelastic collision, conserved quantity is: [1]

- A kinetic energy
- B momentum
- C each body's velocity

2.3 For $k=16$ and $m=4$, angular frequency is _____ rad/s. Show your relation and units. [2]

2.4 Decide whether this claim is true. Justify it: Zero torque about a point implies constant angular momentum _____ about it. [2]

3 Exam-style questions

3.1 A bead of mass m moves without friction along a straight rail. Its potential is $V(x) = ax^4 - bx^2$, with a, b positive. Find every equilibrium, classify stability, and derive the small-oscillation frequency at each stable equilibrium. State why the frequency formula cannot describe motion near $x = 0$. [6]