

5.1 Quantum mechanics and atomic physics

Name: _____ Class: _____ Date: _____

Total: 11 marks

KEY WORDS normalisation • operator

Objective

Use wavefunctions, operators and measurement probabilities; Solve standard bound-state models and tunnelling reasoning; Apply angular momentum, spin and atomic spectral principles. Original public practice; not an ETS paper.

1 Key knowledge and worked examples

Normalise so $\int |\psi|^2 dx = 1$. In an orthonormal expansion $\psi = \sum c_j \varphi_j$ the norm squared is $\sum |c_j|^2$; each energy probability is $|c_j|^2$.

Hermitian observables have real eigenvalues. The expectation $\langle A \rangle = \int \psi^* A \psi dx$ is a repeated-preparation average. Apply operator identities to eigenstates: $A^4 = I$ gives only real eigenvalues $+1$ and -1 .

For an infinite well $0 < x < L$, zero endpoint values give sine modes with $E_n = n^2 \pi^2 \hbar^2 / (2mL^2)$, $n \geq 1$. Doubling L quarters the energies. Finite barriers permit exponential tails.

Commuting observables share an eigenbasis. $T = p^2 / (2m)$ commutes with p ; a variable $V(x)$ generally breaks $[H, p] = 0$. $[J_x, J_y] = i\hbar J_z$; J^2 commutes with each component. Photon energies equal level differences, subject to selection rules.

■ Worked method


An infinite well extends from 0 to L . Vanishing wavefunction at both ends gives

$$\psi_n = \sqrt{2/L} \sin(n\pi x/L), \quad E_n = n^2 \pi^2 \hbar^2 / (2mL^2).$$

Prepare $\psi = (\psi_1 + i\psi_2)/\sqrt{2}$. Orthogonality makes its norm one. Energy measurements give E_1 or $4E_1$ with equal probability:

$$\langle E \rangle = \frac{1}{2}E_1 + \frac{1}{2}(4E_1) = \frac{5}{2}E_1.$$

This average is not an allowed single energy result. The relative phase changes in time, so a superposition of different energies is not a stationary state.



2 Practice

2.1 Choose one answer. In an infinite well, E_3/E_1 is: [1]

- A 3
- B 6
- C 9
- D The boundary conditions determine no energies
- E Replace $\hbar$ with h without changing the coefficient

2.2 Choose one answer. The probability density is: [1]

- A ψ
- B $|\psi|^2$
- C the energy alone
- D The boundary conditions determine no energies
- E Replace $\hbar$ with h without changing the coefficient

2.3 If $E_1 = 2$ eV, $E_2 =$ _____ eV. Show your relation and units. [2]

[2]

[5]