

12.1 Coupled acceleration and relative motion

Name: _____ Class: _____ Date: _____

Total: 11 marks**KEY WORDS** constraint • relative velocity

Objective

Solve coupled-body equations using one acceleration constraint; Transform velocities between inertial frames; Check force and energy limits for constrained motion. Original public practice; not an ETS paper.

1 Key knowledge and worked examples

For table mass m_1 and hanging mass m_2 connected over an ideal pulley, a taut inextensible string gives one acceleration magnitude. Write $T = m_1 a$ and $m_2 g - T = m_2 a$. Thus $a = m_2 g / (m_1 + m_2)$, $T = m_1 m_2 g / (m_1 + m_2) < m_2 g$. Equating tension to weight incorrectly assumes no acceleration.

Equal tensions require an ideal massless string/pulley without friction or inertia. For pulley inertia I , use $(T_2 - T_1)R = I\alpha$ and $a = R\alpha$. Draw separate bodies: a force internal to the combined system remains external to an individual body.

Galilean velocity transformation is $v_{\text{relative}} = v_{\text{ground}} - V$. A load released from a constant-speed level aircraft retains its horizontal ground velocity. With upward positive and no drag, vertical velocity is $-gt$. Relative horizontal velocity is zero; vertical displacement is $-\frac{1}{2}gt^2$, which is a different quantity.

Check limits: $m_1 \rightarrow 0$ gives $a \rightarrow g$ and $T \rightarrow 0$; large m_1 gives $a \rightarrow 0$ and $T \rightarrow m_2 g$ from below. Release-from-rest energy gives $m_2 g s = \frac{1}{2}(m_1 + m_2)v^2$, agreeing with $v^2 = 2as$. This check assumes no friction or other unaccounted work.

■ Worked method

A table mass m_1 and hanging mass m_2 share an acceleration through a taut ideal string. Choose rightward table motion and downward hanging motion positive.

$$\begin{aligned} T &= m_1 a, & m_2 g - T &= m_2 a. \\ a &= \frac{m_2 g}{m_1 + m_2} = \frac{(2 \text{ kg})(10 \text{ m/s}^2)}{3 \text{ kg} + 2 \text{ kg}} = 4 \text{ m/s}^2. \\ T &= m_1 a = (3 \text{ kg})(4 \text{ m/s}^2) = 12 \text{ N}. \end{aligned}$$

T is less than the hanging weight 20 N. A nonzero net downward force is needed for acceleration.

Subtract aircraft velocity; components in m/s

Velocity after 3 s	Horizontal	Vertical
Ground	80	-30
Aircraft	80	0
Payload relative to aircraft	0	-30

2 Practice

2.1 Choose one answer. For $m_1=4$ kg on a frictionless table and $m_2=1$ kg hanging, $g=10$ m/s², tension is: [1]

- A 10 N
- B 2 N
- C 8 N
- D 40 N
- E 0 N

2.2 Choose one answer. A payload is dropped from a constant-speed level aircraft. With $g=10$ m/s² and no drag, its velocity relative to that aircraft after 2 s is: [1]

- A 20 m/s downward
- B 20 m/s forward
- C 40 m/s downward
- D Zero
- E The aircraft speed forward plus 20 m/s downward

2.3 For $m_1=m_2$ and an ideal frictionless table/pulley, $a/g=$ _____ (decimal). Show your relation and units. [2]

