

7.1 Atomic spectra and selection rules

Name: _____ Class: _____ Date: _____

Total: 10 marks

KEY WORDS selection rule • Zeeman splitting

Objective

Relate atomic energies to spectral lines; Apply angular-momentum quantum numbers; Distinguish allowed and forbidden electric-dipole transitions. Original public practice; not an ETS paper.

1 Key knowledge and worked examples

For hydrogen-like one-electron ions, the bound energies scale as $-13.6 Z^2/n^2$ eV in the simplest nonrelativistic model. Emission requires a downward transition and positive photon energy.

Orbital angular momentum has $l=0,\dots,n-1$ and $m_l=-l,\dots,l$. Electron spin adds $s=1/2$; total angular momentum combines orbital and spin contributions.

Electric-dipole selection rules include $\Delta l=\pm 1$ and $\Delta m=0,\pm 1$. These rules concern a specific transition mechanism; forbidden does not mean impossible by every mechanism.

Fine structure, spin-orbit effects and Zeeman splitting refine the simplest hydrogen picture. In many-electron atoms, shielding and electron interactions invalidate direct hydrogen-energy substitution.

■ Worked method

For a nonrelativistic one-electron ion, neglecting fine structure,


$$E_n = -13.6 \text{ eV } Z^2/n^2.$$

For $Z = 2$, an emission from $n = 3$ to $n = 2$ gives positive photon energy:

$$E_\gamma = E_3 - E_2 = (13.6 \text{ eV})(2^2)(1/2^2 - 1/3^2) = 7.56 \text{ eV}.$$

$$\lambda = hc/E_\gamma = (1240 \text{ eV nm})/(7.56 \text{ eV}) = 164 \text{ nm}.$$

An energy difference alone does not guarantee an electric-dipole transition; the participating orbital states must satisfy the selection rules.



2 Practice

2.1 Hydrogen $n=3$ has energy about: [1]

- A. -13.6 eV
B. -1.51 eV
C. $+1.51$ eV

2.2 Which Δl is electric-dipole allowed? [1]

- A. 0
B. 1
C. 2

2.3 For hydrogen $2 \rightarrow 1$, the photon energy is _____ eV. Show your relation and units. [2]

2.4 Decide whether this claim is true. Justify it: Electric-dipole forbidden means no emission mechanism can ever exist. [2]

3 Exam-style questions

3.1 A hydrogenic electron is in a 3d state. Under the electric-dipole approximation, which lower principal-level orbital states can it reach in one emission? Explain why the $n = 1$ level is excluded. Ignore fine structure and spin changes. [4]