

Psychometrics and test construction

GRE Subject Test · GRE Psychology

Scope and prerequisites

GRE Psychology topic 10 (MM.1). Use the overview measurement lesson VI when interpreting research evidence. The practice tasks below are original, not official ETS questions.

Concepts and method

standard error of measurement : The expected spread of observed scores around the true score, $SD \times \sqrt{1-r}$. **construct validity** : Convergent and discriminant evidence that a score reflects the intended trait.

Reliability is consistency, and its type is named by the design that produced it: the same test twice gives test-retest reliability; two similar forms give alternate-forms reliability; consistency within one sitting (split-half, Cronbach alpha) is internal consistency; agreement between scorers is inter-rater reliability. The Spearman-Brown formula $r_{full} = 2r / (1+r)$ rescales a half-test correlation to full test length: $r = .80$ becomes about .89.

Validity is evidence for the proposed interpretation, not of the test itself: content validity asks whether items cover the domain; criterion validity relates scores to an outcome, concurrently or predictively; construct validity assembles convergent and discriminant evidence that the score tracks the intended trait. Reliability caps validity but never establishes it: a consistent scale can systematically measure something else.

Standard scores re-express raw scores against a norm group: $z = (X - \text{mean}) / SD$, $T = 50 + 10z$, and Wechsler deviation $IQ = 100 + 15z$. A z of $+2$ gives IQ 130 and about the 98th percentile. The standard error of measurement $SEM = SD \times \sqrt{1-r}$ converts reliability into a score band: $SD = 15$ with $r = .91$ gives $SEM = 15 \times \sqrt{.09} = 4.5$.

Item analysis checks each item: difficulty p is the proportion answering correctly, and discrimination is the item-total correlation. Items that nearly everyone or no one answers correctly carry little information; negative discrimination flags a likely miskeyed or misleading item.

Choose the reliability coefficient by the source of variation being studied. A standard error of measurement concerns measurement uncertainty under a model; it is not the standard error of a sample mean. The Spearman-Brown estimate assumes sufficiently comparable halves; a formula cannot repair unrelated item content. Standard scores describe relative position and do not show that a test measures the intended construct fairly across groups.

Worked reasoning


Two similar vocabulary tests given a week apart with $r=.90$ show alternate-forms reliability, not test-retest, because the form changed. With $SD=15$ and reliability $.91$, $SEM=4.5$ points. A z of $+2$ corresponds to IQ 130. A split-half correlation of $.80$ corrects to $2 \times .80 / 1.80 \approx .89$ at full length.

A fictional scale has mean 50 points, SD 10 points and reliability $.84$. A score is 65. Compute z and SEM , showing the formulas.


$z = (X - \text{mean}) / SD$; $z = (65 \text{ points} - 50 \text{ points}) / (10 \text{ points}) = 1.5$, dimensionless.
 $SEM = SD \sqrt{1 - r}$; $SEM = (10 \text{ points}) \sqrt{1 - .84} = 4 \text{ points}$. SEM is a model-based uncertainty measure, not proof of validity. $T = 50 + 10z$; $T = 50 + 10(1.5) = 65$ on the T scale.

Location and spread: a limited analogy


Tight
centred



Tight
shifted



Spread
out



Tight grouping is like reliability, being on target like validity: a test can be reliable but not valid

Independent transfer

Two raters repeatedly disagree by five points but rank scripts identically. Does a high correlation establish agreement? Suggest a check.

Attempt independently, then use the matching skill sheet solution.

Review and source limits

Avoid this error: Calling a two-different-forms correlation test-retest reliability, or citing high reliability as if it were validity evidence.