7.1 The covalent model

Name: _____ Class: _____ Date: _____ **Total: 20 marks**

KEY WORDS covalent bond • electronegativity • hydrogen bond • London forces • giant covalent structure • resonance

Objective

Build the skills to answer exam questions on **the covalent model** : Lewis formulas, molecular shape, polarity, giant covalent structures, intermolecular forces and chromatography.

You must be able to:

- draw Lewis formulas and use VSEPR theory to predict shapes and bond angles
- decide whether bonds and molecules are polar from electronegativity and shape
- compare boiling points using London forces, dipole–dipole forces and hydrogen bonds
- describe diamond, graphite, graphene and silicon dioxide, and calculate R_F values

1 Key knowledge and worked examples

Covalent bonds and shapes

A **covalent bond** is the attraction of two nuclei to a shared pair of electrons. In a **coordination bond** both electrons come from one atom (as in NH_4^+).

electron domains	lone pairs	shape	angle	example
2	0	linear	180°	CO_2
3	0	trigonal planar	120°	BF_3
4	0	tetrahedral	109.5°	CH_4
4	1	trigonal pyramidal	107°	NH_3
4	2	bent	104.5°	H_2O

Lone pairs repel more than bonding pairs, so they squeeze the angles. A bond is **polar** when the atoms differ in electronegativity; a molecule is polar only if the bond dipoles do **not** cancel.

intermolecular force	between	strength
London (dispersion)	all molecules; larger with more electrons	weakest
dipole–dipole	polar molecules	stronger
hydrogen bond	H bonded to N, O or F, and a lone pair on N, O or F	strongest

Giant covalent: diamond (each C bonded to four, very hard), graphite (layers, delocalized electrons, conducts), graphene (one layer), SiO_2 (each Si bonded to four O). **Chromatography:** $R_F = \dfrac{\text{distance moved by spot}}{\text{distance moved by solvent}}$.

Worked example 1: shape. SO_2: S has 3 domains with 1 lone pair, so **bent**, a little under 120°.

Worked example 2: boiling points. CH_4 (−162 °C) and C_3H_8 (−42 °C): propane has more electrons, so **stronger London forces**.


Answer shapes: domains, then lone pairs, then shape and angle.

2 Practice


2.1 Deduce the shape and bond angle of an ammonia molecule, NH_3, and of a carbon dioxide molecule, CO_2. [2]



2.2 Both C=O and O−H bonds are polar. Explain why carbon dioxide is a non-polar molecule but water is polar. [3]



2.3 Water boils at 100 °C, but hydrogen sulfide, H_2S, boils at −60 °C. Explain the difference. [3]




exercises.lol exercises.com.cn

3 Exam-style questions

3.1 A student uses paper chromatography to analyse a food dye.

(a) Calculate the R_F value of the spot. [2]




(b) Explain how the components of a mixture are separated. [2]

(c) Explain why the start line is drawn in pencil, not ink. [1]

3.2 Diamond and graphite are both forms of carbon.

(a) Describe the structure of graphite. [2]



(b) Explain why graphite conducts electricity but diamond does not. [2]

(c) Explain why both have very high melting points. [1]

(d) Silicon dioxide also has a very high melting point. Describe its structure. [2]

exercises.lol exercises.com.cn