

<div>T1</div> <div>element</div>	<div>T1</div> <div>compound</div>	<div>T1</div> <div>mixture</div>
<div>T1</div> <div>homogeneous</div>	<div>T1</div> <div>heterogeneous</div>	<div>T1</div> <div>filtration</div>
<div>T1</div> <div>evaporation</div>	<div>T1</div> <div>crystallization</div>	<div>T1</div> <div>distillation</div>

<p>T1</p> <p>A combination of substances that are not chemically bonded together in fixed proportions.</p>	<p>T1</p> <p>A substance containing two or more different elements chemically bonded in fixed proportions.</p>	<p>T1</p> <p>A substance made of atoms with the same proton number.</p>
<p>T1</p> <p>Separating insoluble particles from a fluid by passing the mixture through a porous barrier.</p>	<p>T1</p> <p>Of a mixture: not uniform, with visibly different parts or phases.</p>	<p>T1</p> <p>Of a mixture: uniform throughout, with one phase and no visible boundaries.</p>
<p>T1</p> <p>Separating substances by vaporising and condensing them, using differences in volatility or boiling point.</p>	<p>T1</p> <p>Forming solid crystals from a solution, often by cooling or evaporating the solvent.</p>	<p>T1</p> <p>The escape of particles from the surface of a liquid into its vapour without the whole liquid boiling.</p>

<p>T1</p> <p>solvation</p>	<p>T1</p> <p>saturated</p>	<p>T1</p> <p>kinetic molecular theory</p>
<p>T1</p> <p>sublimation</p>	<p>T1</p> <p>endothermic</p>	<p>T1</p> <p>deposition</p>
<p>T1</p> <p>exothermic</p>	<p>T1</p> <p>purity</p>	<p>T1</p> <p>average kinetic energy</p>

<p>T1</p> <p>The model that matter is made of particles in constant motion whose energy rises with temperature.</p>	<p>T1</p> <p>Describing fatty acids whose hydrocarbon chains contain no carbon-carbon double or triple bonds.</p>	<p>T1</p> <p>Solvent molecules surrounding and attracting dissolved particles.</p>
<p>T1</p> <p>A gas changing directly into a solid.</p>	<p>T1</p> <p>Describing a process that takes in heat from its surroundings.</p>	<p>T1</p> <p>The direct change of a substance from solid to gas without passing through a liquid phase.</p>
<p>T1</p> <p>The mean energy of motion of the particles, proportional to absolute temperature.</p>	<p>T1</p> <p>How free a substance is of other substances; a pure substance has a sharp melting point.</p>	<p>T1</p> <p>Describing a process that transfers heat to its surroundings.</p>

<div>T1</div> <div>absolute zero</div>	<div>T2</div> <div>nucleus</div>	<div>T2</div> <div>proton</div>
<div>T2</div> <div>neutron</div>	<div>T2</div> <div>electron</div>	<div>T2</div> <div>nucleon</div>
<div>T2</div> <div>atomic number</div>	<div>T2</div> <div>mass number</div>	<div>T2</div> <div>nuclear symbol</div>

<p>T2</p> <p>A positively charged subatomic particle found in an atom's nucleus.</p>	<p>T2</p> <p>The small, dense central part of an atom, containing protons and usually neutrons.</p>	<p>T1</p> <p>The lowest temperature on the thermodynamic temperature scale, equal to zero kelvin.</p>
<p>T2</p> <p>A proton or neutron in an atomic nucleus.</p>	<p>T2</p> <p>A negatively charged subatomic particle found outside an atom's nucleus.</p>	<p>T2</p> <p>An electrically neutral subatomic particle found in most atomic nuclei.</p>
<p>T2</p> <p>Notation writing the mass number at the top left and the atomic number at the bottom left of the element symbol.</p>	<p>T2</p> <p>The total number of protons and neutrons in an atomic nucleus.</p>	<p>T2</p> <p>The number of protons in the nucleus of an atom.</p>

<div>T2</div> <div>ion</div>	<div>T2</div> <div>isotope</div>	<div>T2</div> <div>abundance</div>
<div>T3</div> <div>electromagnetic spectrum</div>	<div>T3</div> <div>photon</div>	<div>T3</div> <div>continuous spectrum</div>
<div>T3</div> <div>line emission spectrum</div>	<div>T3</div> <div>energy level</div>	<div>T3</div> <div>excited state</div>

<p>T2</p> <p>The amount or relative frequency of a particular entity, such as a species or an isotope.</p>	<p>T2</p> <p>An atom of an element with the same proton number but a different neutron number from another atom of that element.</p>	<p>T2</p> <p>An atom or group of atoms with a net electric charge from losing or gaining electrons.</p>
<p>T3</p> <p>A spectrum containing all wavelengths in a range without gaps.</p>	<p>T3</p> <p>A quantum of electromagnetic radiation with energy equal to Planck's constant multiplied by its frequency.</p>	<p>T3</p> <p>The complete range of electromagnetic radiation, ordered by frequency or wavelength.</p>
<p>T3</p> <p>States of a system with energy above its ground-state energy.</p>	<p>T3</p> <p>An allowed value of energy that a quantum system can have.</p>	<p>T3</p> <p>Separate coloured lines of light emitted when excited electrons fall to lower levels.</p>

<div>T3</div> <div>quantized</div>	<div>T3</div> <div>converge</div>	<div>T3</div> <div>convergence limit</div>
<div>T3</div> <div>principal quantum number</div>	<div>T3</div> <div>sublevel</div>	<div>T3</div> <div>atomic orbital</div>
<div>T3</div> <div>degenerate</div>	<div>T3</div> <div>Aufbau principle</div>	<div>T3</div> <div>Pauli exclusion principle</div>

<p>T3</p> <p>The frequency where spectral lines merge, used to find ionization energy.</p>	<p>T3</p> <p>Of spectral lines: get closer together toward higher frequency.</p>	<p>T3</p> <p>Able to take only certain fixed values, like the energies of electrons in an atom.</p>
<p>T3</p> <p>A region around the nucleus where an electron is likely to be found; it holds up to two electrons.</p>	<p>T3</p> <p>A division of an energy level: s, p, d or f.</p>	<p>T3</p> <p>The positive integer identifying an electron's main shell and helping determine its orbital energy and size.</p>
<p>T3</p> <p>An orbital holds at most two electrons, and they must have opposite spins.</p>	<p>T3</p> <p>Electrons fill the lowest-energy orbitals available first.</p>	<p>T3</p> <p>Of orbitals: having the same energy, like the three 2p orbitals.</p>

<p>T3</p> <p>opposite spins</p>	<p>T3</p> <p>Hund's rule</p>	<p>T3</p> <p>full electron configuration</p>
<p>T3</p> <p>condensed configuration</p>	<p>T3</p> <p>orbital diagram</p>	<p>T3</p> <p>negative ion</p>
<p>T3</p> <p>positive ion</p>	<p>T3</p> <p>transition element</p>	<p>T4</p> <p>mole</p>

<p>T3</p> <p>The complete list of occupied sublevels, such as $1s^2 2s^2 2p^6 3s^1$ for sodium.</p>	<p>T3</p> <p>Electrons fill degenerate orbitals singly, with parallel spins, before pairing.</p>	<p>T3</p> <p>The two allowed spin states of electrons sharing an orbital, shown as up and down arrows.</p>
<p>T3</p> <p>An atom or group that has gained electrons and carries a negative charge.</p>	<p>T3</p> <p>Boxes for orbitals with arrows for electrons showing their spins.</p>	<p>T3</p> <p>An electron configuration starting from the previous noble gas, such as $[\text{Ne}] 3s^1$.</p>
<p>T4</p> <p>The SI unit of amount of substance, containing exactly 6.02214076 times ten to the power 23 specified entities.</p>	<p>T3</p> <p>A d-block element that forms at least one ion with a partly filled d sublevel.</p>	<p>T3</p> <p>An atom or group that has lost electrons and carries a positive charge.</p>

<p>T4</p> <p>amount of substance</p>	<p>T4</p> <p>Avogadro constant</p>	<p>T4</p> <p>formula unit</p>
<p>T4</p> <p>relative atomic mass</p>	<p>T4</p> <p>relative formula mass</p>	<p>T4</p> <p>molar mass</p>
<p>T4</p> <p>empirical formula</p>	<p>T4</p> <p>molecular formula</p>	<p>T4</p> <p>solution</p>

<p>T4</p> <p>The simplest ratio of ions in an ionic compound, such as NaCl.</p>	<p>T4</p> <p>The number of specified elementary entities in one mole, exactly 6.02214076 times ten to the power 23 per mole.</p>	<p>T4</p> <p>The quantity of particles in a sample, measured in moles.</p>
<p>T4</p> <p>The mass of one mole of a substance.</p>	<p>T4</p> <p>The sum of the relative atomic masses in one formula unit.</p>	<p>T4</p> <p>The weighted mean mass of an element's atoms relative to one twelfth of the mass of a carbon-12 atom.</p>
<p>T4</p> <p>A homogeneous mixture in which one or more solutes are dissolved in a solvent.</p>	<p>T4</p> <p>A formula showing the actual numbers of atoms of each element in one molecule.</p>	<p>T4</p> <p>A formula showing the simplest whole-number ratio of atoms of each element in a compound.</p>

<div>T4</div> <div>solute</div>	<div>T4</div> <div>solvent</div>	<div>T4</div> <div>concentration</div>
<div>T4</div> <div>standard solution</div>	<div>T4</div> <div>Avogadro's law</div>	<div>T5</div> <div>kinetic molecular theory</div>
<div>T5</div> <div>ideal gas</div>	<div>T5</div> <div>elastic</div>	<div>T5</div> <div>pressure</div>

<p>T4</p> <p>The amount of a substance per unit volume of a mixture or solution.</p>	<p>T4</p> <p>The substance in which a solute dissolves to form a solution.</p>	<p>T4</p> <p>A substance dissolved in a solvent to form a solution.</p>
<p>T5</p> <p>The model that matter is made of particles in constant motion whose energy rises with temperature.</p>	<p>T4</p> <p>Equal volumes of all gases, at the same temperature and pressure, contain equal numbers of molecules.</p>	<p>T4</p> <p>A solution of accurately known concentration.</p>
<p>T5</p> <p>Force acting normally per unit area.</p>	<p>T5</p> <p>Of gas collisions: losing no kinetic energy overall.</p>	<p>T5</p> <p>A model of a gas with point particles that exert no intermolecular forces except during perfectly elastic collisions.</p>

<div>T5</div> <div>kelvin scale</div>	<div>T5</div> <div>absolute zero</div>	<div>T5</div> <div>Avogadro's law</div>
<div>T5</div> <div>molar volume</div>	<div>T5</div> <div>ideal gas equation</div>	<div>T5</div> <div>gas constant</div>
<div>T5</div> <div>real gas</div>	<div>T6</div> <div>ionic model</div>	<div>T6</div> <div>ion</div>

<p>T5</p> <p>Equal volumes of all gases, at the same temperature and pressure, contain equal numbers of molecules.</p>	<p>T5</p> <p>The lowest temperature on the thermodynamic temperature scale, equal to zero kelvin.</p>	<p>T5</p> <p>The absolute temperature scale starting at absolute zero; $K = ^\circ C + 273.15$.</p>
<p>T5</p> <p>$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$, the constant in the ideal gas equation.</p>	<p>T5</p> <p>$pV = nRT$, linking pressure, volume, amount and temperature of a gas.</p>	<p>T5</p> <p>The volume of one mole of gas, 22.7 dm^3 at STP.</p>
<p>T6</p> <p>An atom or group of atoms with a net electric charge from losing or gaining electrons.</p>	<p>T6</p> <p>A bonding model in which oppositely charged ions are held together by electrostatic attraction.</p>	<p>T5</p> <p>A gas whose particles have volume and attract each other, deviating at high pressure and low temperature.</p>

T6

cation

T6

anion

T6

transition element

T6

polyatomic ion

T6

ionic bond

T6

electrostatic attraction

T6

electronegativity

T6

lattice

T6

lattice enthalpy

<p>T6</p> <p>A d-block element that forms at least one ion with a partly filled d sublevel.</p>	<p>T6</p> <p>An ion with a negative electric charge.</p>	<p>T6</p> <p>An ion with a positive electric charge.</p>
<p>T6</p> <p>The pull between opposite charges, the basis of ionic and other bonding.</p>	<p>T6</p> <p>The electrostatic attraction between oppositely charged ions.</p>	<p>T6</p> <p>A group of covalently bonded atoms that carries an overall charge.</p>
<p>T6</p> <p>The energy to separate one mole of an ionic solid into gaseous ions.</p>	<p>T6</p> <p>A regular, repeating arrangement of particles or points in a solid.</p>	<p>T6</p> <p>The tendency of an atom in a chemical bond to attract the shared electrons towards itself.</p>

<p>T6</p> <p>volatility</p>	<p>T7</p> <p>covalent bond</p>	<p>T7</p> <p>intermolecular force</p>
<p>T7</p> <p>shared pair</p>	<p>T7</p> <p>octet rule</p>	<p>T7</p> <p>incomplete octet</p>
<p>T7</p> <p>lone pair</p>	<p>T7</p> <p>Lewis formula</p>	<p>T7</p> <p>coordination bond</p>

T7	T7	T6
An attraction between separate molecules.	A chemical bond involving a shared pair of electrons attracted to the nuclei of bonded atoms.	The tendency of a substance to vaporise readily under given conditions.
T7	T7	T7
An atom with fewer than eight outer electrons in a molecule, such as boron in BF_3 .	Atoms tend to bond until they have eight electrons in their outer shell.	Two electrons shared between two atoms, forming a covalent bond.
T7	T7	T7
A covalent bond in which both shared electrons come from one atom.	A diagram showing all valence electrons as bonding pairs and lone pairs.	A pair of outer-shell electrons on one atom that is not shared in a covalent bond.

<p>T7</p> <p>valence shell electron pair repulsion</p>	<p>T7</p> <p>electron domain</p>	<p>T7</p> <p>molecular geometry</p>
<p>T7</p> <p>electronegativity</p>	<p>T7</p> <p>polar bond</p>	<p>T7</p> <p>polar</p>
<p>T7</p> <p>dipole moment</p>	<p>T7</p> <p>covalent network structure</p>	<p>T7</p> <p>allotrope</p>

T7	T7	T7
The 3-D arrangement of the atoms in a molecule.	A region of electron density around an atom: a lone pair or a single, double or triple bond.	The theory that electron domains around a central atom repel and spread as far apart as possible.
T7	T7	T7
Having a separation of charge, with a partly positive and a partly negative end.	A covalent bond with unequal sharing because the atoms differ in electronegativity.	The tendency of an atom in a chemical bond to attract the shared electrons towards itself.
T7	T7	T7
A different structural form of the same element, such as diamond and graphite.	A giant structure of atoms joined by covalent bonds, such as diamond or silicon dioxide.	A measure of the charge separation in a bond or molecule.

<p>T7</p> <p>delocalized</p>	<p>T7</p> <p>silicon dioxide</p>	<p>T7</p> <p>London force</p>
<p>T7</p> <p>dipole–dipole force</p>	<p>T7</p> <p>hydrogen bonding</p>	<p>T7</p> <p>van der Waals force</p>
<p>T7</p> <p>volatility</p>	<p>T7</p> <p>chromatography</p>	<p>T7</p> <p>stationary phase</p>

T7	T7	T7
A weak attraction from temporary induced dipoles, present between all molecules.	The compound SiO_2 , whose common solid forms contain a network of silicon and oxygen atoms joined by covalent bonds.	Of electrons: spread over several atoms instead of held between two.
T7	T7	T7
A general name for the weak intermolecular forces: London and dipole forces.	An attraction involving a hydrogen atom bonded to an electronegative atom and an electron-rich site on another atom.	An attraction between the permanent dipoles of polar molecules.
T7	T7	T7
The phase that remains fixed in a chromatography system and interacts differently with components of a mixture.	A separation technique based on different affinities of substances for a stationary phase and a mobile phase.	The tendency of a substance to vaporise readily under given conditions.

<div>T7</div> <div>mobile phase</div>	<div>T7</div> <div>retardation factor</div>	<div>T8</div> <div>metallic model</div>
<div>T8</div> <div>cation</div>	<div>T8</div> <div>lattice</div>	<div>T8</div> <div>delocalized electron</div>
<div>T8</div> <div>metallic bond</div>	<div>T8</div> <div>non-directional</div>	<div>T8</div> <div>electrical conductivity</div>

<p>T8</p> <p>A bonding model with positive metal ions held together by attraction to delocalized electrons.</p>	<p>T7</p> <p>Rf: distance moved by a spot divided by distance moved by the solvent front.</p>	<p>T7</p> <p>The solvent that moves through the stationary phase in chromatography.</p>
<p>T8</p> <p>An electron distributed over several atoms rather than confined to one atom or a single bond.</p>	<p>T8</p> <p>A regular, repeating arrangement of particles or points in a solid.</p>	<p>T8</p> <p>An ion with a positive electric charge.</p>
<p>T8</p> <p>A measure of how readily a material allows electric current to flow through it.</p>	<p>T8</p> <p>Acting equally in all directions, like metallic and ionic bonding.</p>	<p>T8</p> <p>The electrostatic attraction between a lattice of cations and the delocalized electrons.</p>

<div>T8</div> <div>thermal conductivity</div>	<div>T8</div> <div>malleability</div>	<div>T8</div> <div>ductility</div>
<div>T9</div> <div>material</div>	<div>T9</div> <div>bonding triangle</div>	<div>T9</div> <div>alloy</div>
<div>T9</div> <div>polymer</div>	<div>T9</div> <div>monomer</div>	<div>T9</div> <div>repeating unit</div>

<p>T8</p> <p>the strain at fracture, the property heat treatment trades away for strength</p>	<p>T8</p> <p>The ability to be hammered into shape without breaking.</p>	<p>T8</p> <p>Delocalized electrons (and vibrating ions) pass kinetic energy quickly through the metal.</p>
<p>T9</p> <p>A mixture of a metal with other metals, or with a non-metal such as carbon.</p>	<p>T9</p> <p>A triangle chart using electronegativity to place a substance between ionic, covalent and metallic bonding.</p>	<p>T9</p> <p>A substance chosen for its properties to make something, such as a metal, ceramic, polymer or composite.</p>
<p>T9</p> <p>The part of a polymer chain that repeats again and again.</p>	<p>T9</p> <p>A molecule that can join with other molecules to form a polymer.</p>	<p>T9</p> <p>A large molecule made from many linked repeating units derived from smaller molecules.</p>

<div>T9</div> <div>biodegradable</div>	<div>T9</div> <div>addition polymerization</div>	<div>T10</div> <div>periodic table</div>
<div>T10</div> <div>periodicity</div>	<div>T10</div> <div>period</div>	<div>T10</div> <div>group</div>
<div>T10</div> <div>block</div>	<div>T10</div> <div>metalloid</div>	<div>T10</div> <div>alkali metal</div>

<p>T10</p> <p>An arrangement of chemical elements by atomic number that displays recurring patterns in their properties.</p>	<p>T9</p> <p>Joining monomers with C=C double bonds into a polymer with no other product.</p>	<p>T9</p> <p>Able to be broken down by living organisms.</p>
<p>T10</p> <p>A vertical column in the periodic table whose elements have related outer-electron configurations and chemical properties.</p>	<p>T10</p> <p>A horizontal row of the periodic table, corresponding to the principal shell occupied by the outer electrons of its elements.</p>	<p>T10</p> <p>A repeating pattern in element properties across the periodic table.</p>
<p>T10</p> <p>A group 1 metal, such as sodium, that reacts vigorously with water.</p>	<p>T10</p> <p>An element with properties between metals and non-metals, such as silicon.</p>	<p>T10</p> <p>The region of the periodic table named by the sublevel being filled: s, p, d or f.</p>

<div>T10</div> <div>halogen</div>	<div>T10</div> <div>nuclear charge</div>	<div>T10</div> <div>shielding</div>
<div>T10</div> <div>electron affinity</div>	<div>T10</div> <div>basic</div>	<div>T10</div> <div>acidic</div>
<div>T10</div> <div>amphoteric</div>	<div>T10</div> <div>acid deposition</div>	<div>T10</div> <div>ocean acidification</div>

<p>T10</p> <p>The reduction in nuclear attraction experienced by an electron because of repulsion from other electrons.</p>	<p>T10</p> <p>The total positive charge of an atomic nucleus, determined by its number of protons.</p>	<p>T10</p> <p>The elements in group 17 of the periodic table.</p>
<p>T10</p> <p>Of an oxide: reacting with bases, typical of non-metal oxides.</p>	<p>T10</p> <p>Of an oxide: reacting with acids, typical of metal oxides.</p>	<p>T10</p> <p>The energy change when a gaseous atom gains one electron.</p>
<p>T10</p> <p>The fall in ocean pH as seawater absorbs more carbon dioxide.</p>	<p>T10</p> <p>Acidic substances from sulfur and nitrogen oxides falling to the ground as rain or dry particles.</p>	<p>T10</p> <p>Able to react as both an acid and a base, like aluminium oxide.</p>

<p>T11</p> <p>The C_6H_5- group, a benzene ring attached to the rest of a molecule.</p>	<p>T11</p> <p>A formula drawn as lines, with carbon at corners and line ends and hydrogens on carbon not shown.</p>	<p>T10</p> <p>The charge an atom would have if all its bonds were ionic.</p>
<p>T11</p> <p>A compound with the same molecular formula but atoms joined in a different order.</p>	<p>T11</p> <p>A family of compounds with the same functional group and the same general formula.</p>	<p>T11</p> <p>An atom or group of atoms responsible for characteristic chemical reactions of an organic compound.</p>
<p>T11</p> <p>Of an alcohol or halogenoalkane: the functional carbon bonded to three other carbons.</p>	<p>T11</p> <p>Of an alcohol or halogenoalkane: the functional carbon bonded to two other carbons.</p>	<p>T11</p> <p>Of an alcohol or halogenoalkane: the functional carbon bonded to one other carbon.</p>

<p>T11</p> <p>index of hydrogen deficiency</p>	<p>T11</p> <p>mass spectrometer</p>	<p>T11</p> <p>molecular ion</p>
<p>T11</p> <p>wavenumber</p>	<p>T11</p> <p>fingerprint region</p>	<p>T11</p> <p>nuclear magnetic resonance</p>
<p>T11</p> <p>chemical shift</p>	<p>T11</p> <p>integration</p>	<p>T12</p> <p>exothermic</p>

T11	T11	T11
An ion formed from a whole molecule by gaining or losing electrons without breaking the molecule into fragments.	An instrument that separates ions by mass-to-charge ratio to find isotopes and their abundances.	The number of rings plus π bonds in a molecule, found from its formula.
T11	T11	T11
A technique using magnetic fields to show the environments of hydrogen atoms in a molecule.	The complex infrared region below about 1500 cm^{-1} , unique to each compound.	The reciprocal of wavelength, in cm^{-1} , used for infrared absorptions.
T12	T11	T11
Describing a process that transfers heat to its surroundings.	Measuring the area under an NMR signal to determine the relative number of nuclei producing it.	The position of an NMR signal relative to a reference, reflecting the chemical environment of the nucleus.

<div>T12</div> <div>enthalpy change</div>	<div>T12</div> <div>endothermic</div>	<div>T12</div> <div>activation energy</div>
<div>T12</div> <div>limiting reactant</div>	<div>T12</div> <div>uncertainty</div>	<div>T13</div> <div>energy cycle</div>
<div>T13</div> <div>bond enthalpy</div>	<div>T13</div> <div>average bond enthalpy</div>	<div>T13</div> <div>Hess's law</div>

<p>T12</p> <p>The minimum energy that reacting particles need for a successful reaction.</p>	<p>T12</p> <p>Describing a process that takes in heat from its surroundings.</p>	<p>T12</p> <p>The heat transferred in a reaction or process at constant pressure.</p>
<p>T13</p> <p>A diagram of different routes between the same start and end, used with Hess's law.</p>	<p>T12</p> <p>An estimate of the possible range of error in a measurement.</p>	<p>T12</p> <p>The reactant that is used up first and therefore limits the amount of product formed.</p>
<p>T13</p> <p>The total enthalpy change of a reaction is the same whatever route is taken, because energy is conserved.</p>	<p>T13</p> <p>The mean energy to break one mole of a bond, averaged over many compounds.</p>	<p>T13</p> <p>The energy needed to break one mole of a covalent bond, with the substances in the gaseous state.</p>

<div>T14</div> <div>functional unit</div>	<div>T14</div> <div>life-cycle assessment</div>	<div>T15</div> <div>conservation of mass</div>
<div>T15</div> <div>coefficient</div>	<div>T15</div> <div>mole ratio</div>	<div>T15</div> <div>stoichiometry</div>
<div>T15</div> <div>molar volume</div>	<div>T15</div> <div>titration</div>	<div>T15</div> <div>standard solution</div>

<p>T15</p> <p>The principle that total mass remains unchanged in an ordinary chemical reaction within a closed system.</p>	<p>T14</p> <p>Assessing a product's environmental impact from raw materials to disposal.</p>	<p>T14</p> <p>A specified amount of service or function used as the common basis for comparing products in a life-cycle assessment.</p>
<p>T15</p> <p>The quantitative relationships between reactants and products specified by a balanced chemical equation.</p>	<p>T15</p> <p>The ratio of amounts of substances measured in moles, often given by the coefficients in a balanced equation.</p>	<p>T15</p> <p>A numerical or constant factor multiplying a term in an expression.</p>
<p>T15</p> <p>A solution of accurately known concentration.</p>	<p>T15</p> <p>A method of finding an unknown concentration by measuring the volume needed to react with a solution of known concentration.</p>	<p>T15</p> <p>The volume of one mole of gas, 22.7 dm³ at STP.</p>

<p>T15</p> <p>pipette</p>	<p>T15</p> <p>indicator</p>	<p>T15</p> <p>burette</p>
<p>T15</p> <p>end point</p>	<p>T15</p> <p>titre</p>	<p>T15</p> <p>concordant</p>
<p>T15</p> <p>limiting reactant</p>	<p>T15</p> <p>in excess</p>	<p>T15</p> <p>theoretical yield</p>

T15	T15	T15
A graduated tube with a tap used to deliver accurately measured volumes of liquid.	A substance that shows a change in chemical conditions through an observable change, often in colour.	Glassware that delivers an exact volume of liquid.
T15	T15	T15
Of titres: agreeing closely, usually within 0.10 cm^3 .	The volume of solution added from the burette to reach the end point.	The observed signal used to stop a titration, such as an indicator's colour change, which should be close to equivalence.
T15	T15	T15
The mass of product calculated from the limiting reactant.	Present in more than the amount needed to react completely.	The reactant that is used up first and therefore limits the amount of product formed.

<div>T15</div> <div>atom economy</div>	<div>T16</div> <div>rate of reaction</div>	<div>T16</div> <div>tangent</div>
<div>T16</div> <div>initial rate</div>	<div>T16</div> <div>collision theory</div>	<div>T16</div> <div>activation energy</div>
<div>T16</div> <div>Maxwell–Boltzmann distribution</div>	<div>T16</div> <div>surface area</div>	<div>T16</div> <div>catalyst</div>

<p>T16</p> <p>A straight line that touches a curve at a point and has the same direction there.</p>	<p>T16</p> <p>The change in the amount or concentration of a reactant or product per unit time.</p>	<p>T15</p> <p>The mass of the desired product as a percentage of the total mass of all products.</p>
<p>T16</p> <p>The minimum energy that reacting particles need for a successful reaction.</p>	<p>T16</p> <p>The explanation that reactions require collisions with sufficient energy and suitable orientation.</p>	<p>T16</p> <p>The rate of a reaction or process measured at or very near its beginning.</p>
<p>T16</p> <p>A substance that increases a reaction rate without being used up overall.</p>	<p>T16</p> <p>The total area of the outer surfaces of a three-dimensional object.</p>	<p>T16</p> <p>The graph of how many particles have each kinetic energy at a given temperature.</p>

<p>T16</p> <p>alternative reaction pathway</p>	<p>T17</p> <p>reversible reaction</p>	<p>T17</p> <p>dynamic equilibrium</p>
<p>T17</p> <p>closed system</p>	<p>T17</p> <p>isotope</p>	<p>T17</p> <p>equilibrium constant</p>
<p>T17</p> <p>position of equilibrium</p>	<p>T17</p> <p>Le Chatelier's principle</p>	<p>T17</p> <p>yield</p>

<p>T17</p> <p>A state in which opposing processes continue at equal rates, so there is no overall change in composition.</p>	<p>T17</p> <p>A reaction whose products can react to form the original reactants under suitable conditions.</p>	<p>T16</p> <p>A different route with lower activation energy, provided by a catalyst.</p>
<p>T17</p> <p>A value relating the equilibrium activities, concentrations or partial pressures of products and reactants at a specified temperature.</p>	<p>T17</p> <p>An atom of an element with the same proton number but a different neutron number from another atom of that element.</p>	<p>T17</p> <p>A system that does not exchange matter with its surroundings, although energy may be transferred.</p>
<p>T17</p> <p>The amount of product obtained in a reaction, often expressed as a percentage of the theoretical amount.</p>	<p>T17</p> <p>When a system at equilibrium is disturbed, the position of equilibrium shifts to reduce the effect of the change.</p>	<p>T17</p> <p>The relative amounts of reactants and products present when forward and reverse reaction rates are equal.</p>

<p>T17</p> <p>compromise</p>	<p>T18</p> <p>proton</p>	<p>T18</p> <p>Brønsted–Lowry acid</p>
<p>T18</p> <p>Brønsted–Lowry base</p>	<p>T18</p> <p>conjugate acid–base pair</p>	<p>T18</p> <p>amphiprotic</p>
<p>T18</p> <p>ionic product of water</p>	<p>T18</p> <p>strong acid</p>	<p>T18</p> <p>weak acid</p>

T18	T18	T17
A proton (H^+) donor.	A positively charged subatomic particle found in an atom's nucleus.	Industrial conditions chosen to balance yield, rate and cost.
T18	T18	T18
Able both to donate and to accept a proton, like water.	Two species differing by exactly one proton, such as NH_4^+ and NH_3 .	A proton (H^+) acceptor.
T18	T18	T18
An acid that only partially ionises in aqueous solution.	An acid that essentially completely ionises in aqueous solution at the concentrations being considered.	$K_w = [\text{H}^+][\text{OH}^-] = 1.00 \times 10^{-14}$ at 298 K.

<p>T18</p> <p>neutralization</p>	<p>T18</p> <p>pH curve</p>	<p>T18</p> <p>equivalence point</p>
<p>T19</p> <p>oxidation</p>	<p>T19</p> <p>reduction</p>	<p>T19</p> <p>oxidation state</p>
<p>T19</p> <p>oxidizing agent</p>	<p>T19</p> <p>reducing agent</p>	<p>T19</p> <p>half-equation</p>

<p>T18</p> <p>The stage of a titration at which the reactants have been mixed in their stoichiometric proportions.</p>	<p>T18</p> <p>A graph of pH against volume added during a titration.</p>	<p>T18</p> <p>The reaction of an acid with a base to form a salt and water.</p>
<p>T19</p> <p>The charge an atom would have if all its bonds were ionic.</p>	<p>T19</p> <p>A chemical change involving a gain of electrons or a decrease in oxidation state.</p>	<p>T19</p> <p>A chemical change involving a loss of electrons or an increase in oxidation state.</p>
<p>T19</p> <p>An equation showing just the oxidation or just the reduction, with electrons.</p>	<p>T19</p> <p>A substance that donates electrons to another substance and is itself oxidised.</p>	<p>T19</p> <p>The one that is reduced.</p>

<p>T19</p> <p>activity series</p>	<p>T19</p> <p>displacement</p>	<p>T19</p> <p>primary cell</p>
<p>T19</p> <p>anode</p>	<p>T19</p> <p>cathode</p>	<p>T19</p> <p>salt bridge</p>
<p>T19</p> <p>secondary cell</p>	<p>T19</p> <p>fuel cell</p>	<p>T19</p> <p>electrolytic cell</p>

T19	T19	T19
A cell that cannot be recharged.	A reaction in which one element replaces another in a compound.	A list of metals in order of how readily they lose electrons.
T19	T19	T19
A link of ions between two half-cells that completes the circuit.	An electrode at which reduction occurs.	An electrode at which oxidation occurs.
T19	T19	T19
A cell in which an external electrical supply drives a non-spontaneous chemical reaction.	An electrochemical device that converts the chemical energy of a continuously supplied fuel and oxidant into electricity.	A rechargeable cell.

<div>T19</div> <div>electrolysis</div>	<div>T19</div> <div>electrolyte</div>	<div>T19</div> <div>distillation</div>
<div>T19</div> <div>reflux</div>	<div>T19</div> <div>hydrogenation</div>	<div>T20</div> <div>free radical</div>
<div>T20</div> <div>unpaired electron</div>	<div>T20</div> <div>homolytic fission</div>	<div>T20</div> <div>single-barbed arrow</div>

<p>T19</p> <p>Separating substances by vaporising and condensing them, using differences in volatility or boiling point.</p>	<p>T19</p> <p>A substance that conducts electricity when molten or dissolved because it contains mobile ions.</p>	<p>T19</p> <p>The decomposition of an electrolyte by an electric current passing through its molten form or a solution.</p>
<p>T20</p> <p>A species containing one or more unpaired electrons.</p>	<p>T19</p> <p>The addition of hydrogen to a substance, often across a carbon-carbon double bond using a catalyst.</p>	<p>T19</p> <p>Heating a reaction mixture while condensing its vapour and returning the condensate to the same vessel.</p>
<p>T20</p> <p>A half-headed arrow showing the movement of one electron.</p>	<p>T20</p> <p>Breaking a covalent bond so that each fragment receives one electron from the bonding pair.</p>	<p>T20</p> <p>An electron alone in an orbital, as in a free radical.</p>

<div>T20</div> <div>ultraviolet light</div>	<div>T20</div> <div>non-polar</div>	<div>T20</div> <div>radical substitution</div>
<div>T20</div> <div>chain reaction</div>	<div>T20</div> <div>initiation</div>	<div>T20</div> <div>propagation</div>
<div>T20</div> <div>termination</div>	<div>T20</div> <div>product mixture</div>	<div>T20</div> <div>chlorofluorocarbon</div>

<p>T20</p> <p>Substitution by a radical chain: initiation by UV, propagation steps, and termination when radicals join.</p>	<p>T20</p> <p>Having no overall separation of charge.</p>	<p>T20</p> <p>High-energy light that can split bonds to form radicals.</p>
<p>T20</p> <p>A chain step in a radical reaction that consumes one radical and produces another.</p>	<p>T20</p> <p>The first stage of a chain reaction, in which reactive species such as free radicals are produced.</p>	<p>T20</p> <p>A reaction in which each step makes a radical that continues the chain.</p>
<p>T20</p> <p>A compound of carbon, chlorine and fluorine that releases chlorine radicals in the stratosphere.</p>	<p>T20</p> <p>The mixture of several products formed, as in radical substitution.</p>	<p>T20</p> <p>The ending stage of a process, such as completion of protein synthesis or the stopping of a reaction chain.</p>

<div>T20</div> <div>stratosphere</div>	<div>T20</div> <div>ozone</div>	<div>T21</div> <div>nucleophile</div>
<div>T21</div> <div>electrophile</div>	<div>T21</div> <div>heterolytic fission</div>	<div>T21</div> <div>curly arrow</div>
<div>T21</div> <div>halogenoalkane</div>	<div>T21</div> <div>nucleophilic substitution</div>	<div>T21</div> <div>leaving group</div>

<p>T21</p> <p>An electron-pair donor that forms a covalent bond to an electron-deficient centre.</p>	<p>T20</p> <p>O₃, which absorbs harmful ultraviolet light in the stratosphere.</p>	<p>T20</p> <p>The atmosphere layer from about 10 to 50 km up, containing the ozone layer.</p>
<p>T21</p> <p>An arrow showing the movement of an electron pair in a mechanism.</p>	<p>T21</p> <p>Breaking a covalent bond so that both electrons in the bonding pair go to the same fragment.</p>	<p>T21</p> <p>A species that accepts an electron pair to form a covalent bond.</p>
<p>T21</p> <p>The atom or group that departs with the bonding pair in a substitution.</p>	<p>T21</p> <p>A reaction in which an electron-pair donor replaces a leaving group in a molecule.</p>	<p>T21</p> <p>An alkane derivative in which one or more hydrogen atoms have been replaced by halogen atoms.</p>

<div>T21</div> <div>electrophilic addition</div>	<div>T22</div> <div>systematic error</div>	<div>T22</div> <div>uncertainty</div>

<p>T22</p> <p>An estimate of the possible range of error in a measurement.</p>	<p>T22</p> <p>A consistent measurement bias caused by the method or instrument.</p>	<p>T21</p> <p>A reaction in which an electron-pair acceptor initiates addition across a multiple bond.</p>