

Introduction to the particulate nature of matter · supported focus

International Baccalaureate · IB Diploma · Chemistry · SL

Scope and prerequisites

Supported SL focus. First assessment 2025; current brief acquired; full Chemistry guide not acquired. Remaining guide, assessment and practical requirements retain their recorded holds.

Prerequisites: read the stated quantities and units, use arithmetic and the model conditions below. Each lesson develops its own method before independent transfer.

These are original or explicitly fictional teaching examples, not actual measurements or completed assessed learner investigations.


Gas models and absolute temperature

■ What would explain this observation?

- A sealed gas container changes pressure when heated. Celsius ratios cannot predict the pressure change because the gas model uses **absolute temperature**.
- Start with a prediction. State the quantities or features you would compare, then decide what evidence could distinguish two explanations.

■ Build the model

- In a gas model, particles move randomly and pressure results from momentum transfer at walls. The **ideal gas** equation connects pressure, volume, amount and absolute temperature.
- absolute temperature: Temperature on the kelvin scale; ideal gas: A gas model with specified simplifying assumptions.



■ Choose evidence that can test it

- At fixed amount and volume, pressure is proportional to kelvin temperature. At fixed temperature and amount, pressure is inversely proportional to volume. State which quantities are fixed before choosing a relationship.
 - Use approved apparatus with a temperature range and pressure limit set by the teacher. Allow thermal equilibrium and record pressure against kelvin temperature. Never heat an improvised sealed vessel.
-

■ Work from known quantities

- State the known values and their units. Choose the relation because its assumptions fit this case, then rearrange before substitution.
- Known: pressure is 100 kPa at 300 K, with fixed volume and amount. At 330 K, $p_2/p_1 = T_2/T_1$. $p_2 = p_1 \cdot T_2/T_1 = 100 \times 330/300 = 110$ kPa. A 30 °C rise is a 30 K change, but the temperature ratio must use kelvin.

Example:

At fixed volume, pressure is 120 kPa at 300 K. Find pressure at 350 K. Use the same sequence: known quantities → model → relation → substitution → unit and interpretation.

■ Check the conclusion and its limits

- An ideal gas is a model with conditions of validity. Celsius zero is not zero molecular motion, and internal energy is not determined by pressure alone.
- Return to the original observation. Explain what the result supports, which conditions it assumes, and one way to test a competing explanation.

Warn:

A gas pressure ratio can always use Celsius temperatures. This claim is false: An ideal gas is a model with conditions of validity. Celsius zero is not zero molecular motion, and internal energy is not determined by pressure alone.

Key:

Gas models and absolute temperature: At fixed amount and volume, pressure is proportional to kelvin temperature. At fixed temperature and amount, pressure is inversely proportional to volume. State which quantities are fixed before choosing a relationship.