

Course Companion

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Every topic handout in one volume

exercises.lol

Contents

1. Foundation	3
2. Ecology	5
3. Biodiversity and conservation	7
4. Water	9
5. Land	11
6. Atmosphere and climate change	13
7. Natural resources	15
8. Human populations and urban systems	18
9. Environmental law	20
10. Environmental and ecological economics	23

11. Environmental ethics.....	26
12. Experimental programme.....	29

Foundation

International Baccalaureate · IB Diploma · Environmental Systems
and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Environmental systems, feedback and perspectives

■ What would explain this observation?

- A lake can respond to the same nutrient input differently depending on its current state. Environmental outcomes depend on **stocks**, **flows** and feedback.
-

■ Build the model

- A system model has a boundary, stores and flows.
- Negative feedback counteracts change; positive feedback amplifies it.
- An environmental perspective shapes what people value and which management outcomes they prioritize.
- stock : An amount held within a system.
- flow : Transfer of an amount per unit time.



$$\text{Final stock} = \text{initial stock} + \text{inflows} - \text{outflows} = 800 + 120 - 90 = 830$$

■ Choose evidence that can test it

- Separate physical evidence from value judgements.
- A stock is an amount at a time; a flow is an amount per time.
- Sustainability decisions require ecological, social and economic evidence and a stated scale.

In the lab: Draw the system boundary before calculating a budget. Identify inflows, outflows and possible unmeasured terms. Compare stakeholder claims using the same evidence, then explain where values lead to different choices.

■ Work from known quantities

- Known: a reservoir begins with 800 units, receives 120 and loses 90 during a week.
- Change in storage = inflow-outflow = $120 - 90 = 30$.
- Final storage = $800 + 30 = 830$ units.
- The model assumes no omitted flow.

Your turn: A stock starts at 600, receives 80 and loses 110. Find final stock.

■ Check the conclusion and its limits

- A balance that closes mathematically can still omit a real process if terms were estimated to fit.
- Positive feedback does not mean a socially desirable result.

Not true: Positive feedback always produces a socially beneficial outcome. “Positive” in positive feedback means the change is AMPLIFIED, not that the outcome is good. Melting permafrost releasing methane is positive feedback with harmful consequences; desirability is a separate judgement.

Key:

- A system model has a boundary, stores and flows.
- Separate physical evidence from value judgements.
- Watch for: A balance that closes mathematically can still omit a real process if terms were estimated to fit.

■ Write from a final stock of 475

- A store starts at 450 units, receives 60 and loses 35.
- Find the final stock. Positive feedback means a change is amplified, not that the outcome is good.

Ecology

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Sampling populations without choosing the answer

■ What would explain this observation?

- A field edge looks richer in plants than its centre. Choosing only the richest patches would build the desired result into the sampling method.
-

■ Build the model

- A **population** consists of organisms of one species in a defined area.
- Random **quadrats** estimate density without deliberately selecting patches.
- A transect investigates change along an environmental gradient.
- quadrat : A defined area used for sampling.
- population: Organisms of one species in a defined area.



Quadrat positions chosen before looking at the plants avoid sampling bias

■ Choose evidence that can test it

- Estimate total abundance by multiplying mean density by area, with consistent units.
- This assumes sampled areas represent the habitat.
- Patchiness and too few samples widen uncertainty.

In the lab: Choose coordinates with random numbers before visiting the patches. Record quadrat area and counting rules. For a transect, use fixed distances and measure a relevant abiotic variable. Do not damage habitats or sample unsafe locations.

■ Work from known quantities

- Known: five 0.25 square metre quadrats contain 3, 4, 6, 5 and 2 plants.
- Mean count = $20/5 = 4$.
- Density = $4/0.25 = 16$ plants per square metre.
- Estimated abundance in 100 square metres = $16 \times 100 = 1,600$ plants.

Your turn: Mean count is 6 in a 0.5 square metre quadrat. Find density per square metre.

■ Check the conclusion and its limits

- Quadrats suit organisms that do not move quickly.
- A food chain arrow shows the direction of energy transfer, not the direction a predator travels.

Not true: Sampling only the most crowded patches gives an unbiased population estimate. Choosing the most crowded patches overestimates density, because the samples are not representative of the whole area. Unbiased estimation needs random or systematic sampling decided before looking at the plants.

Key:

- A population consists of organisms of one species in a defined area.
- Estimate total abundance by multiplying mean density by area, with consistent units.
- Watch for: quadrats suit organisms that do not move quickly.

■ Write from a density of 20 plants per square metre

- The mean count is 5 plants in a 0.25 square metre quadrat.
- Find the density per square metre. Sampling only crowded patches overestimates density.

Biodiversity and conservation

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Biodiversity, conservation and sampling evidence

■ What would explain this observation?

- A site with many individuals can still be dominated by one species. Abundance, richness and **evenness** describe different features of biodiversity.

■ Build the model

- **Species richness** counts species.
- Evenness concerns relative abundance.
- Conservation can protect habitats, populations or genetic diversity; a management judgement must state the conservation goal.
- species richness: The number of species recorded.
- evenness: How evenly individuals are distributed among species.



Same richness, different evenness: site B is more even

■ Choose evidence that can test it

- Compare surveys with similar area, effort, season and identification rules.
- A diversity index is meaningful only with its formula and conventions specified.
- Habitat fragmentation can affect movement and gene flow even when total area changes little.

In the lab: Use non-destructive field sampling approved by the school. Identify organisms with a suitable key and record uncertain identifications rather than inventing species. Combine ecological evidence with stakeholder perspectives on land use.

■ Work from known quantities

- Known: site A has counts 8, 1 and 1; site B has 4, 3 and 3.
- Both have richness 3 and total abundance 10.
- The most abundant species occupies 80% in A and 40% in B.
- B is more even under these counts.

Your turn: A survey records 6, 5, 3 and 2 individuals in four species. Find species richness.

■ Check the conclusion and its limits

- An index cannot by itself explain the cause of a difference.
- A newly recorded species may reflect improved observation rather than a recent ecological arrival.

Not true: A larger total number of individuals always means greater species richness. Richness counts how many species are present, not how many individuals. Ten thousand individuals of one species give richness 1, while ten species with one individual each give richness 10.

Key:

- Species richness counts species.
- Compare surveys with similar area, effort, season and identification rules.
- Watch for: an index cannot by itself explain the cause of a difference.

■ Write from a richness of 5

- A survey records 7, 2, 2, 2 and 1 individuals across five species.
- State the species richness. A larger total of individuals is not greater richness.

Water

International Baccalaureate · IB Diploma · Environmental Systems
and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Water, pollution and management trade-offs

■ What would explain this observation?

- A downstream sample has less dissolved oxygen after a nutrient input. The explanation must connect biological processes with the water-system evidence.
-

■ Build the model

- **Eutrophication** can begin with nutrient enrichment, followed by greater producer growth and increased decomposition after organisms die.
- Decomposer respiration can lower dissolved oxygen.
- Water availability depends on quantity, quality and access.
- eutrophication: Nutrient enrichment with ecological consequences.
- **pollutant load** : Mass of pollutant transferred per time.



$$Load = concentration \times flow = 2 \text{ mg/L} \times 500 \text{ L/min} = 1,000 \text{ mg/min} = 1 \text{ g/min}$$

■ Choose evidence that can test it

- Distinguish concentration from total pollutant load.
- Dilution can reduce concentration without removing the pollutant mass.
- A management plan should consider upstream causes, users and effects on other parts of the catchment.

In the lab: Collect approved water-quality observations at matched sites and times. Use calibrated instruments and consistent sampling depth. Avoid contact with contaminated water and do not infer potability from clarity.

■ Work from known quantities

- Known: concentration is 2 mg per litre and flow is 500 litres per minute.
- Load = concentration $\times$ flow = $2 \times 500 = 1,000$ mg per minute = 1 g per minute.
- A lower concentration during high flow can still produce a high total load.

Your turn: Concentration is 3 mg/L and flow 200 L/min. Find load in mg/min.

■ Check the conclusion and its limits

- Clear water is not necessarily safe water.
- Nutrient enrichment does not directly use up oxygen; the pathway includes biological activity and decomposition.

Not true: Clear water is always safe to drink. Clarity shows only that little is suspended. Dissolved metals, pathogens and many chemicals are invisible; safety requires testing against standards, not looking at the water.

Key:

- Eutrophication can begin with nutrient enrichment, followed by greater producer growth and increased decomposition after organisms die.
- Distinguish concentration from total pollutant load.
- Watch for: clear water is not necessarily safe water.

■ Write from a load of 400 mg per minute

- Concentration is 5 mg per litre and flow is 80 litres per minute.
- Find the load. Clear water is not evidence that the water is safe to drink.

Land

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Land, food and resource decisions

■ What would explain this observation?

- A farm can increase yield while increasing soil loss or water demand. Food-system evaluation requires more than tonnes harvested.
-

■ Build the model

- Soil supports nutrient cycling, water storage and organisms.
- **Erosion** removes soil; nutrient depletion changes fertility.
- **Resource** use involves extraction, production, distribution and waste.
- erosion: Removal and transport of soil or rock.
- resource: A material or service used to meet needs.



Compare yield per hectare, not total harvest: $18 \text{ t} \div 6 \text{ ha} = 3 \text{ t/ha}$ against $20 \text{ t} \div 10 \text{ ha} = 2 \text{ t/ha}$

■ Choose evidence that can test it

- Distinguish total yield from yield per area and energy input.
- Sustainable management must consider regeneration rates and impacts beyond the farm boundary.
- Different stakeholders may rank food security, biodiversity and income differently.

In the lab: Compare management options using a stated functional unit and the same timescale. Use approved secondary datasets or non-destructive soil observations. Explain trade-offs instead of claiming a method has no disadvantages.

■ Work from known quantities

- Known: farm A harvests 18 tonnes from 6 hectares.
- Yield per area = $18/6 = 3$ tonnes per hectare.
- Farm B harvests 20 tonnes from 10 hectares, so its yield is 2 tonnes per hectare despite a larger total harvest.

Your turn: A farm harvests 28 tonnes from 7 hectares. Find yield per hectare.

■ Check the conclusion and its limits

- A high yield is not proof of long-term soil health.
- Renewable resources can be depleted when use exceeds regeneration.

Not true: A renewable resource cannot be depleted. Renewable means the resource can regenerate, not that it regenerates faster than every possible rate of use. Fish stocks and soils are renewable yet have been depleted when harvest exceeded renewal.

Key:

- Soil supports nutrient cycling, water storage and organisms.
- Distinguish total yield from yield per area and energy input.
- Watch for: A high yield is not proof of long-term soil health.

■ Write from a yield of 5 tonnes per hectare

- A farm harvests 40 tonnes from 8 hectares.
- Find the yield per hectare. A renewable resource can still be depleted.

Atmosphere and climate change

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Climate evidence, energy budgets and policy

■ What would explain this observation?

- One cold day does not disprove a warming climate. Weather describes short-term conditions; climate describes distributions over longer times and regions.

■ Build the model

- The Earth energy balance includes incoming solar radiation, reflection, absorption and outgoing infrared radiation.
- Greenhouse gases absorb and emit infrared radiation.
- Feedback can alter the response to an initial forcing.
- **mitigation** : Actions addressing causes of environmental change.
- **adaptation** : Actions reducing harm from environmental impacts.



$$\text{Absorbed} = \text{incoming} - \text{reflected}: 200 - 50 = 150 \text{ units}$$

■ Choose evidence that can test it

- Distinguish mitigation, which addresses drivers, from adaptation, which reduces harm from impacts.
- A policy assessment needs evidence about effectiveness, cost, equity and uncertainty; one criterion is not the entire decision.

In the lab: Compare multi-year data using consistent baselines. State the region, timescale and uncertainty. At HL, connect a management decision to law, economics and ethics rather than treating these lenses as extra definitions only.

■ Work from known quantities

- Known: a surface receives 200 power units and reflects 50.
- Absorbed input = incoming-reflected = $200 - 50 = 150$.
- Reflected fraction = $50/200 = 0.25 = 25\%$.
- A change in reflectivity alters the absorbed budget under this model.

Your turn: Incoming energy is 240 units and reflected energy 72. Find reflected percentage.

■ Check the conclusion and its limits

- The greenhouse effect is not the same process as ozone depletion.
- A carbon footprint estimate depends on its system boundary.

Not true: One local weather observation establishes a global climate trend. Weather is a short-term local state; climate is the long-term statistical pattern. One cold day in one city cannot confirm or refute a global trend built from decades of worldwide records.

Key:

- The Earth energy balance includes incoming solar radiation, reflection, absorption and outgoing infrared radiation.
- Distinguish mitigation, which addresses drivers, from adaptation, which reduces harm from impacts.
- Watch for: the greenhouse effect is not the same process as ozone depletion.

■ Write from 20 percent reflected

- A surface receives 500 energy units and reflects 100.
- Find the reflected percentage and the absorbed input. One local weather observation does not establish a climate trend.

Natural resources

International Baccalaureate · IB Diploma · Environmental Systems
and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Natural resources: soils, yields and trade-offs

■ What would explain this observation?

- Two farms report the same total harvest. The one using half the land is not automatically more sustainable: **natural resources** regenerate at their own rates.
-

■ Build the model

- Soil is a resource: its removal and transport, by water, wind or gravity, is a loss measured beyond one season.
- Sustainable management weighs total **yield** per area and per energy input against regeneration rates and impacts beyond the farm boundary.
- Different stakeholders may rank food security, biodiversity and income differently.
- yield: The useful output of a managed system, per stated basis.
- natural resources: Materials and energy sources supplied by ecosystems and used by people.



■ Choose evidence that can test it

- Compare management options using a stated **functional unit** and the same timescale.
- Use approved secondary datasets or non-destructive soil observations.
- Explain trade-offs: no method is without disadvantages.

In the lab: State the functional unit (per hectare, per tonne, per gigajoule) before comparing, and keep the timescale identical.

■ Work from known quantities

- Known: farm A harvests 12 t from 4 ha (3 t per ha) using 40 GJ; farm B harvests 12 t from 2 ha (6 t per ha) using 25 GJ.

Your turn: A third farm matches B's per-hectare yield but feeds its soil nothing back. Which basis of comparison is still missing?

■ Check the conclusion and its limits

- A comparison is only as good as its stated functional unit and timescale.
- Stakeholder priorities differ: a food-security ministry and a conservation body can rank the same data differently, both correctly.

Not true: The higher-yielding farm is automatically the more sustainable one: re-generation and beyond-boundary impacts decide otherwise.

Key:

- State the functional unit and the timescale before comparing.
- Total yield, yield per area and per energy input are different claims.
- Watch for: impacts beyond the farm boundary and differing stakeholder priorities.

■ **Write from a functional unit**

- Two farms are compared per farm and per 1 000 kg of grain.
- Per farm compares whole farms; per 1 000 kg of grain compares equal food output. Choose a functional unit that matches the stated question, with equivalent function and the same timescale.

■ **Write from three yield claims**

- A farm yields 12 t on 2 ha with 40 GJ of energy input.
- Total 12 t; per area $12 \div 2 = 6$ t per ha; per energy $12 \div 40 = 0.3$ t per GJ – three different claims from one farm.

■ **Write from a boundary stakeholder**

- Irrigation water leaves the farm boundary and reaches downstream users.
- Name them in the trade-off: an impact beyond the boundary is still an impact of the system.

Human populations and urban systems

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Population, urban systems and indicators

■ What would explain this observation?

- A city can grow because people move into it even when births equal deaths. Population change has several flows.

■ Build the model

- Population change depends on births, deaths, **immigration** and emigration.
- Urban systems exchange energy, materials and waste with a wider region.
- A sustainability **indicator** measures a specified feature rather than total wellbeing.
- immigration: Movement into a population.
- indicator: A measure of a specified condition.



$$\text{Change} = \text{births} - \text{deaths} + \text{immigration} - \text{emigration} = +150 \text{ people, } 1.5\% \text{ of } 10,000$$

■ Choose evidence that can test it

- Compare rates with a common denominator and interval.
- An average can hide unequal access to services.
- At HL, evaluate how economic incentives, legal responsibilities and ethical priorities change an urban management proposal.

In the lab: Use published demographic and city-service data with dates and definitions. Protect privacy when collecting local opinions. Compare the same geographical boundary rather than mixing an urban core with a whole metropolitan region.

■ Work from known quantities

- Known: a city has 10,000 people; births 150, deaths 100, immigration 300 and emigration 200 in one year.
- Change = $150 - 100 + 300 - 200 = 150$.
- Growth percentage = $150 / 10,000 \times 100 = 1.5\%$.

Your turn: A population of 2,000 grows by 40 in a year. Find percentage growth.

■ Check the conclusion and its limits

- A projected population is conditional on assumptions.
- Urban growth does not automatically mean improved health, income or environmental quality for every resident.

Not true: Population growth depends only on births and deaths. Population change = births - deaths + immigration - emigration. A country with more deaths than births can still grow through net immigration, so births and deaths alone do not decide it.

Key:

- Population change depends on births, deaths, immigration and emigration.
- Compare rates with a common denominator and interval.
- Watch for: A projected population is conditional on assumptions.

■ Write from 1 percent growth

- A population of 5,000 has 90 births, 40 deaths, 30 immigrants and 30 emigrants in one year.
- Find the percentage growth. Population change is not births and deaths alone.

Environmental law

International Baccalaureate · IB Diploma · Environmental Systems
and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Environmental law: rules, evidence and enforcement

■ What would explain this observation?

- Two catchments have the same discharge limit but different water quality. A legal rule can influence behaviour only through its coverage, implementation and **enforcement**.
-

■ Build the model

- Environmental law defines obligations and permissions within a **jurisdiction**.
- A discharge permit may specify a concentration, a total load, monitoring and consequences of non-compliance.
- International agreements require implementation through institutions; signing an agreement alone does not show that an environmental outcome has improved.
- enforcement: Implementation of rules through monitoring and consequences.
- jurisdiction: The geographical or institutional scope of an authority.



A concentration limit is not a load limit: the same 1.5 mg/L at double the flow doubles the load

■ Choose evidence that can test it

- Distinguish the legal instrument from the evidence about its effect.
- A concentration limit can be satisfied while total pollutant load rises if flow increases.
- Evaluate monitoring coverage, detection probability, incentives, access to justice and effects on different groups.
- An observed change after a policy is compatible with an effect but also with rainfall, industrial changes or other interventions.

In the lab: Use an attributed historical or clearly fictional permit and paired monitoring data. Identify the regulated activity, responsible authority, geographical boundary and date. Compare upstream/downstream and before/after observations where available, document sampling gaps, and discuss whose evidence or costs are excluded. Do not present this classroom model as advice about current local law.

■ Work from known quantities

- Known: a fictional permit caps discharge at 2 mg/L.
- A monitored discharge is 1.5 mg/L at 400 L/min.
- Load = concentration $\times$ flow = 600 mg/min.
- At the same concentration and 800 L/min the load is 1,200 mg/min.
- Both concentration observations satisfy the fictional cap, while the second transports twice the mass.

Your turn: A discharge contains 1.2 mg/L at 500 L/min. Calculate transported mass per minute.

■ Check the conclusion and its limits

- Compliance with one indicator does not prove ecological safety or fair enforcement.
- Compare actual outcomes with the policy objective and consider both administrative feasibility and affected communities.

Not true: A concentration-compliant discharge necessarily has a small total pollutant load. Concentration compliance limits how strong the discharge is, not how much total mass flows. Load = concentration x flow, so a compliant concentration at very high flow can deliver a large pollutant load.

Key:

- Environmental law defines obligations and permissions within a jurisdiction.
- Distinguish the legal instrument from the evidence about its effect.
- Watch for: compliance with one indicator does not prove ecological safety or fair enforcement.

■ Write from a load of 800 mg per minute

- A discharge is 0.8 mg per litre at 1,000 litres per minute.
- Find the load. Meeting a concentration limit does not prove the total load is small.

■ Write from a lower concentration but a higher load

- A second discharge is 0.5 mg per litre but flows at 2,400 litres per minute.
- Find the load: $0.5 \times 2,400 = 1,200$ mg per minute – the cleaner-looking discharge carries the bigger load.

■ Write from a daily load

- The first discharge (0.8 mg per litre at 1,000 litres per minute) runs all day.
- Find the daily load: $800 \times 60 \times 24 = 1,152,000$ mg – about 1.2 kg per day for one pipe.

Environmental and ecological economics

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Environmental economics: costs beyond the market price

■ What would explain this observation?

- A cheap product can impose waste or pollution costs on people who did not buy it. Its market price may omit part of its environmental cost.

■ Build the model

- An **externality** is a cost or benefit affecting a third party that is not fully reflected in a transaction.
- Environmental economics examines incentives and valuation, including instruments such as taxes, subsidies and tradable permits.
- Ecological economics also questions growth assumptions and emphasizes ecological limits and relationships between the economy and its supporting systems.
- externality: A transaction effect on a third party not fully reflected in its price.
- **social cost** : Private cost plus external cost within the stated model.



$$\text{Social cost} = \text{private cost} + \text{external cost} = 1,600 + 600 = 2,200 \text{ units}$$

■ Choose evidence that can test it

- In a simplified model, marginal social cost equals marginal private cost plus marginal external cost.
- A pollution charge can alter incentives, but selecting a charge requires evidence and a stated objective.
- Monetary totals do not by themselves settle distribution, irreversibility, uncertainty or whether a service can be substituted.

In the lab: Construct a transparent table with a common currency, time period and system boundary. Separate measured expenditure from estimated environmental damage. Compare at least two valuation assumptions, identify who pays and who benefits, and consider a non-monetary indicator such as habitat condition. Label these as classroom scenarios rather than forecasts.

■ Work from known quantities

- Known: producing one item costs the producer 8 units, while the scenario estimates an external cost of 3 units.
- Social cost is 11 units per item.
- For 200 items, private cost is 1,600 units and estimated external cost is 600 units, giving 2,200 units total under this model.
- If external damage is estimated at 1–5 units per item, the social-cost range is 1,800–2,600 units.

Your turn: Private cost is 6 units per item and external cost is 2 units. Calculate social cost for 50 items.

■ Check the conclusion and its limits

- A money value is an assumption-dependent estimate, not a measurement of intrinsic ecological value.
- A policy that lowers total cost can still place a disproportionate burden on a vulnerable group.

Not true: The cheapest market price necessarily identifies the smallest total environmental cost. Market price reflects private costs. External costs - pollution, health damage - fall on others and are absent from the price, so the cheapest market option can carry the largest total cost.

Key:

- An externality is a cost or benefit affecting a third party that is not fully reflected in a transaction.

- In a simplified model, marginal social cost equals marginal private cost plus marginal external cost.
- Watch for: A money value is an assumption-dependent estimate, not a measurement of intrinsic ecological value.

■ **Write from a social cost of 900**

- Private cost is 5 units per item and the estimated external cost is 4 units. The scenario covers 100 items.
- Find the total social cost. The cheapest market price can omit external costs.

■ **Write from a corrective tax**

- The same external cost (4 per item) is collected as a tax on the producer.
- What must the market price now carry? Private 5 + tax 4 = 9 – the full social cost moves into the price.

■ **Write from 1,000 of social cost**

- A second scenario has 200 items, private cost 3 each and external cost 2 each.
- Find the total social cost: $200 \times (3 + 2) = 1,000$ – externality times quantity scales the gap.

Environmental ethics

International Baccalaureate · IB Diploma · Environmental Systems
and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Environmental ethics: make the value judgement explicit

■ What would explain this observation?

- People may agree on the measured loss of a wetland but disagree on whether a development is acceptable. The disagreement can concern values as well as scientific uncertainty.
-

■ Build the model

- Environmental ethics examines reasons for responsibilities toward people, other organisms and ecological systems.
- Anthropocentric reasoning emphasizes human interests; biocentric reasoning recognizes moral standing of living organisms; ecocentric reasoning emphasizes ecological wholes.
- These positions are lenses for analysing arguments, and real individuals can combine them.
- **intrinsic value** : Value considered independent of usefulness to another party.
- **environmental justice** : Fair treatment in environmental burdens, benefits and decision-making.



Measure the change first, then argue from a stated value position

■ Choose evidence that can test it

- Separate descriptive claims from normative claims.
- A species count is evidence about the sampled community; deciding that every species has intrinsic value is a value position.
- Compare consequences, rights, duties and fairness across generations.
- A defensible argument states its values, uses accurate evidence and acknowledges competing claims rather than disguising a value choice as a calculation.

In the lab: Use a fictional planning dispute or an attributed historical case. Build an argument table: stakeholder, factual claim, evidence, value principle and possible response. Include affected communities and future generations without pretending to speak for them. Use voluntary anonymous classroom positions; assess the reasoning rather than agreement with a preferred worldview.

■ Work from known quantities

- Known: a wetland proposal affects 12 hectares of a 60-hectare site, or 20%.
- This quantifies the stated area change.
- An anthropocentric argument might examine flood protection and livelihoods; a biocentric argument might consider affected organisms; an ecocentric argument might consider hydrological connections.
- The 20% figure cannot alone decide which ethical argument is strongest.

Your turn: A project affects 9 hectares within a 45-hectare site. Calculate the percentage of area affected.

■ Check the conclusion and its limits

- Worldview labels are analytical aids, not fixed stereotypes of cultures or people.
- Neither a stakeholder majority nor a numerical index automatically resolves a rights-based or intergenerational objection.

Not true: A percentage of habitat lost alone settles the ethical acceptability of a development. The 20% is a measurement; acceptability is a judgement that also weighs what the habitat means, who is affected and what alternatives exist. Different ethical frameworks can weigh the same 20% differently.

Key:

- Environmental ethics examines reasons for responsibilities toward people, other organisms and ecological systems.
- Separate descriptive claims from normative claims.
- Watch for: worldview labels are analytical aids, not fixed stereotypes of cultures or people.

■ Write from 30 percent of the site

- A proposal affects 15 hectares of a 50-hectare site.
- Find the percentage of area affected. That percentage does not settle whether the proposal is ethically acceptable.

Experimental programme

International Baccalaureate · IB Diploma · Environmental Systems and Societies · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2026; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Uncertainty, gradients and model testing

■ What would explain this observation?

- A line passing near every data point is useful, but its gradient can still be uncertain. A graph is evidence for a model within the measurement range.

■ Build the model

- Random variation makes repeated readings differ.
- **Systematic error** shifts results consistently.
- Absolute **uncertainty** has the measured unit; relative or percentage uncertainty compares uncertainty with the measured value.
- uncertainty: A quantified limitation on a measured result.
- systematic error: A consistent measurement bias.



Every point lies within its uncertainty of the line of best fit

■ Choose evidence that can test it

- For a product or quotient, adding fractional uncertainties is a common maximum-uncertainty approximation.
- For a difference, add absolute uncertainties.
- A nonzero intercept can reveal an offset or an incomplete model.

In the lab: Show units on axes and choose a sensible scale. Plot uncertainty bars where justified, draw a best-fit line rather than joining every point, and estimate steepest and shallowest plausible gradients when the course method calls for them.

■ Work from known quantities

- Known: length = 50.0 mm with uncertainty 1.0 mm.
- Percentage uncertainty = $\text{absolute uncertainty} / \text{value} \times 100 = 1.0 / 50.0 \times 100 = 2.0\%$.
- For a quotient of two independently measured quantities with maximum percentage uncertainties 2% and 3%, the summed maximum estimate is 5%.

Your turn: A 40 cm reading has an absolute uncertainty of 1 cm. Find percentage uncertainty.

■ Check the conclusion and its limits

- Repeating readings reduces random uncertainty in a mean but does not automatically remove a zero error.
- Do not quote more decimal places than your measurement can support.

Not true: Repeating a measurement always removes a calibration offset. Repeating and averaging reduces RANDOM scatter in the mean. A calibration offset shifts every reading identically, so repeats land on the same wrong value; only recalibration removes it.

Key:

- Random variation makes repeated readings differ.
- For a product or quotient, adding fractional uncertainties is a common maximum-uncertainty approximation.
- Watch for: repeating readings reduces random uncertainty in a mean but does not automatically remove a zero error.

■ Write from a percentage uncertainty of 5 percent

- A reading is 80 cm and the absolute uncertainty is 4 cm.
- Find the percentage uncertainty. Repeating a measurement does not remove a calibration offset.