3.5 Vectors: lines and the scalar product

Name: _____ Class: _____ Date: _____ **Total: 28 marks**

KEY WORDS vector • magnitude • scalar product • position vector • skew lines • velocity

Objective

Build the skills to answer exam questions on **vectors** : magnitude, the scalar product, angles, vector equations of lines, intersecting and skew lines, and motion.

You must be able to:

- find magnitudes, unit vectors and position vectors
- use the scalar product to find angles and test for perpendicular vectors
- write vector equations of lines and decide whether two lines intersect, are parallel or are skew
- model motion with constant velocity, $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$

1 Key knowledge and worked examples

■ Vectors and lines

$\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$; $|\mathbf{v}| = \sqrt{v_1^2 + v_2^2 + v_3^2}$; a unit vector is $\dfrac{\mathbf{v}}{|\mathbf{v}|}$.

Scalar product: $\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3 = |\mathbf{a}||\mathbf{b}| \cos\theta$. Perpendicular $\Leftrightarrow \mathbf{a} \cdot \mathbf{b} = 0$.

exercises.lol exercises.com.cn

Line: $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$ (a point plus multiples of a direction). For two lines:

test	result
directions are multiples	parallel (or the same line)
equations for x, y, z all agree for one λ, μ	intersect
not parallel and the equations do not all agree	**skew**


Motion: position $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$; speed $= |\mathbf{v}|$.

Worked example. $\mathbf{a} = \begin{pmatrix}1\\2\\2\end{pmatrix}$, $\mathbf{b} = \begin{pmatrix}3\\0\\4\end{pmatrix}$: $\mathbf{a} \cdot \mathbf{b} = 11$, $|\mathbf{a}| = 3$, $|\mathbf{b}| = 5$, $\cos\theta = \frac{11}{15}$, $\theta = \mathbf{42.8}°$.


Answer shapes: for intersection, solve two components and **check the third**.

2 Practice

2.1 Find the magnitude of $\mathbf{v} = \begin{pmatrix}2\\-1\\2\end{pmatrix}$ and a unit vector in its direction. [2]



2.2 Find the angle between $\begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 4 \\ 8 \end{pmatrix}$. [3]




2.3 Find p such that $\begin{pmatrix} p \\ 2 \\ -1 \end{pmatrix}$ is perpendicular to $\begin{pmatrix} 3 \\ p \\ 4 \end{pmatrix}$. [3]



exercises.lol exercises.com.cn

3 Exam-style questions

3.1 $L_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$ and $L_2 : \mathbf{r} = \begin{pmatrix} 3 \\ -1 \\ 4 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}$.




(a) Show that L_1 and L_2 are not parallel. [1]

(b) Show that L_1 and L_2 are skew. [4]

(c) Find the acute angle between the directions of L_1 and L_2. [3]

(d) Find the point on L_1 where $x = 7$. [2]

3.2 Boat A has position $\begin{pmatrix}2\\1\end{pmatrix}+t\begin{pmatrix}3\\4\end{pmatrix}$ and boat B has position $\begin{pmatrix}10\\-3\end{pmatrix}+t\begin{pmatrix}-1\\6\end{pmatrix}$, in km, t hours after noon.



(a) Find the speed of boat A. [2]

(b) Show that the boats collide, and find where. [4]

(c) Find the angle between their paths. [4]

exercises.lol exercises.com.cn