

Exercise walk-through

4.1

Data summaries, histograms and interpretation

International Baccalaureate · IB Diploma · Mathematics: Applications and Interpretation

4.1 Data summaries, histograms and interpretation

Name: _____ Class: _____ Date: _____ **Total: 13 marks**

KEY WORDS frequency density

Objective

Data summaries, histograms and interpretation. Original classroom practice for IB HL; not an official IB paper.

1 Key knowledge and worked examples

■ Can one average tell the whole story?

- Two groups have the same median but different spread. One summary cannot describe both location and consistency.
- This lesson studies **frequency density** : Frequency divided by class width,

used as histogram height.

■ The method


- Compare an appropriate average and spread in context.
 - A histogram uses area for frequency, so height=frequency/class width.
 - Grouped estimates assume representative values within intervals.
-

■ A worked case

$$\bar{x} = \frac{\sum x_i}{n}, \quad \text{density} = \frac{\text{frequency}}{\text{class width}}$$

Example:

A class from 10 to 20 with frequency 30 has density $30/10=3$. A class from 20 to 40 with frequency 20 has density $20/20=1$. Its wider bar must not be mistaken for a larger density. For values 2,4,4,6,9, the median is 4 and mean is 5.



■ Where it goes wrong

- The tallest histogram bar need not contain the most observations.
- A grouped mean is an estimate.
- Correlation does not prove causation, and extrapolation extends beyond the observed range.

Not true: A histogram bar's height always equals its frequency, even with unequal class widths. With unequal class widths the height is frequency density; frequency is the area of the bar.

■ Applying it

- Choose a display that fits the data type.
- Give both a numerical comparison and what it means for the population; do not infer more precision than the sample supports.

■ In your course

- Applications and Interpretation HL, the course first assessed in 2021.
- Every paper allows a calculator, but method marks still need the expression or equation written before its value.

Key:

- **Frequency density:** Frequency divided by class width, used as histogram height.
 - Compare an appropriate average and spread in context.
 - Watch for: the tallest histogram bar need not contain the most observations.
-

■ **Write from a mean of 9**

- A class of width 2 has frequency 22.
- The values 5, 8, 8, 11 and 13 have a mean and a median. Bar height is not always the frequency.

Exercises 2.1-2.3

2.1 Find the density for frequency 30 and class width 10. Show the relation, substitution and units. [2]

2.2 Find the mean of 2,4,4,6,9. Show the relation, substitution and units. [2]

2.3 Find the median of 2,4,4,6,9. Show the relation, substitution and units. [2]

Exercises 2.1-2.3

2.1 Find the density for frequency 30 and class width 10. Show the relation, substitution and units. [2]

2.1 $\text{Density} = \text{frequency} / \text{class width} = 30 / 10 = 3.$

2.2 Find the mean of 2,4,4,6,9. Show the relation, substitution and units. [2]

2.2 The total is 25 across five observations: $\text{mean} = 25 / 5 = 5.$

2.3 Find the median of 2,4,4,6,9. Show the relation, substitution and units. [2]

2.3 The sorted list is 2,4,4,6,9. Its third observation is 4.

Exercises 2.4-3.1

2.4 Decide whether this claim is true and justify: Frequency density means Frequency divided by class width, used as histogram height. [2]

2.5 In a histogram with unequal class widths, which bar contains the most observations? Choose one answer.

- **A** Always the widest bar, because a wider class must contain more values
- **B** The one with the largest AREA, not necessarily the tallest
- **C** Always the tallest bar, because height shows how many observations it holds [1]

3.1 Classes 0–10 and 10–30 have frequencies 20 and 30. Find histogram densities and compare bar heights with frequencies; can the exact mean be recovered? [4]

Exercises 2.4-3.1

2.4 Decide whether this claim is true and justify: Frequency density means Frequency divided by class width, used as histogram height. [2]

2.4 True. True: Frequency divided by class width, used as histogram height.

2.5 In a histogram with unequal class widths, which bar contains the most observations? Choose one answer.

- **A** Always the widest bar, because a wider class must contain more values
- **B** The one with the largest AREA, not necessarily the tallest
- **C** Always the tallest bar, because height shows how many observations it holds [1]

2.5 B Histogram area equals frequency. A wide bar of low density can hold more observations than a narrow tall bar; height alone shows density, not frequency.

3.1 Classes 0–10 and 10–30 have frequencies 20 and 30. Find histogram densities and compare bar heights with frequencies; can the exact mean be recovered? [4]

3.1 Densities 2 and 1.5, so the higher bar has fewer observations. Areas represent frequency. Only a grouped midpoint estimate, not the exact mean, follows without raw values; state the endpoint convention.