5.5 Differential equations and numerical solutions

Name: _____ Class: _____ Date: _____ **Total: 13 marks**

KEY WORDS initial condition

Objective

Differential equations and numerical solutions. Original classroom practice for IB HL; not an official IB paper.

1 Key knowledge and worked examples

What does the starting value decide?

- A growing population's rate is proportional to its current size. The rate equation describes a whole family until an initial population is supplied.
- This lesson studies **initial condition** : A specified value that selects a particular solution from a family.

The method


- For dy/dx=ky, separate variables: dy/y=k dx, giving y=Ae^(kx).
- Use an initial condition to find A.
- Euler's method takes y next=y+h f(x,y), with a chosen step h.

A worked case

$$\frac{dy}{dx} = ky, \qquad y = Ae^{kx}$$

Example:

With dy/dx=0.5y and y(0)=2, y=2e^(0.5x). Euler with h=0.2 gives y(0.2)≈2+0.2×1=2.2 and y(0.4)≈2.2+0.2×1.1=2.42. The exact second value is about 2.4428.



Where it goes wrong

- An initial condition determines the integration constant; it is not a replacement for integrating.
- Dividing by y can omit an equilibrium solution y=0.
- Euler accuracy depends on step size and the equation.

Not true: Every step size gives exactly the same Euler approximation. Euler's method depends on the step size; smaller steps usually give a more accurate value.

Applying it

- For second-order linear equations, combine the complementary function with an appropriate particular integral, then apply the required initial conditions.
- This is Further Pure scope; check the named unit.

In your course

- Applications and Interpretation HL, the course first assessed in 2021.
- Every paper allows a calculator, but method marks still need the expression or equation written before its value.

Key:

- **Initial condition**: A specified value that selects a particular solution from a family.
- For dy/dx=ky, separate variables: dy/y=k dx, giving y=Ae^(kx).
- Watch for: an initial condition determines the integration constant; it is not a replacement for integrating.

■ **Write from a first Euler value of 6.6**

- Use Euler with step 0.2 for y prime = 0.5 y, starting at y = 6.
- Find the first two approximations. A different step size gives a different approximation.

2 Practice

2.1 Using Euler with h=0.2 for y prime=0.5y,y_0=2, find y_1. Show the relation, substitution and units. [2]

2.2 For that approximation, find y_2. Show the relation, substitution and units. [2]

2.3 For y=2e^(0.5x), find y at x=0. Show the relation, substitution and units. [2]

2.4 Decide whether this claim is true and justify: Every step size gives exactly the same Euler approximation. [2]

2.5 For dy/dx=0.5y with y(0)=2, what does the initial condition determine? Choose one answer.

- **A** The step size for Euler's method, which is set equal to y(0)=2
- **B** The growth constant k, which changes from 0.5 to 2 once y(0)=2 is used
- **C** The constant A in the family y=Ae^(0.5x), giving y=2e^(0.5x) [1]

3 Exam-style questions

3.1 Use Euler h=0.5 for y′=−y, y(0)=2 to estimate y(1). Compare with 2 e^−1 and state a limitation. [4]

exercises.lol exercises.com.cn