

Functions

International Baccalaureate · IB Diploma · Mathematics:
Applications and Interpretation · SL

Scope and prerequisites

This handout covers selected SL topics for the course first assessed in 2021; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.

Domains, inverses and composition

■ Which inputs are allowed?

- A square-root model returns a real output only for some inputs. Its formula alone does not specify a complete function.
 - This lesson studies **domain** : The set of allowed inputs to a function.
-

■ The method


- State the domain and range.
 - For an inverse, first ensure the function is one-to-one on its domain.
 - Composition fg means apply g first, then f ; the intermediate output must be an allowed input to f .
-

■ A worked case

$$f(g(x)) = (f \circ g)(x), \quad f^{-1}(f(x)) = x$$

Example:

For $f(x)=\sqrt{x-2}$, $x \geq 2$ and the range is $y \geq 0$. From $y=\sqrt{x-2}$, $x=y^2+2$. Thus f inverse(x)= x^2+2 with $x \geq 0$. For $g(x)=x+3$, $fg(1)=f(4)=\sqrt{2}$.



$y = \sqrt{x - 2}$ exists only for $x \geq 2$, so its domain starts at 2 and its range starts at 0

■ Where it goes wrong

- Squaring can introduce extraneous solutions.
- Restricting a parabola's domain is essential before claiming an inverse.
- A horizontal translation inside f has the opposite sign to the graph's movement.

Not true: Every quadratic function on all real numbers has an inverse function. A quadratic on all real numbers is many-to-one ($f(1)=f(-1)$ for x^2), so its domain must be restricted first.

■ Applying it

- Check $f(f^{-1}(x))=x$ on the inverse domain.
 - Use a sketch to test whether a horizontal line meets the original graph more than once.
-

■ In your course

- Applications and Interpretation SL, the course first assessed in 2021.
- Every paper allows a calculator, but method marks still need the expression or equation written before its value.

Key:

- **Domain:** The set of allowed inputs to a function.

- State the domain and range.
- Watch for: squaring can introduce extraneous solutions.

■ Write from an inverse value of 7

- The function $4x + 1$ sends 7 to 29.
- Find the input that the inverse sends 29 to. A quadratic needs a restricted domain.

Regression, financial models and residuals

■ Is the prediction supported by the data?

- A line predicts bus travel time from distance. A small calculation error matters less than using a model outside its evidence.
 - This lesson studies **residual** : The observed value minus the value predicted by a fitted model.
-

■ The method


- A linear model $y=a+bx$ has intercept a and slope b with contextual units.
 - Inspect residuals and the data range.
 - For repeated percentage change use a geometric model; for a loan distinguish principal, rate, repayment and period.
-

■ A worked case

$$\hat{t} = 8 + 2d, \quad e = t - \hat{t}$$

Example:

The fitted model $t=8+2d$ predicts 18 minutes at $d=5$ km. If the observed time is 21, the residual is 3 minutes. A ¥1000 deposit at 5% compound annual interest becomes $1000 \times 1.05^3 = 1157.625$ after 3 years.



$$\text{Residual} = \text{observed} - \text{predicted} = 21 - 18 = 3 \text{ minutes}$$

■ Where it goes wrong

- A good fit does not prove a causal mechanism.
- A correlation coefficient measures linear association, not the gradient.
- Calculator output must be translated into a model, checked and interpreted.

Not true: A high correlation permits reliable extrapolation to any distance. A model fitted to data is reliable only near that data; extrapolating far outside it is unreliable.

■ Applying it

- In IB AI, record the data source, domain, assumptions and calculator method.
- Compare an alternative model and judge predictions against residuals.
- A financial answer must state payment timing and whether interest is compounded.

■ In your course

- Applications and Interpretation SL, the course first assessed in 2021.

Key:

- **Residual:** The observed value minus the value predicted by a fitted model.
- A linear model $y = a + bx$ has intercept a and slope b with contextual units.
- Watch for: A good fit does not prove a causal mechanism.

■ Write from a prediction of 17

- The model $t = 5 + 3d$ predicts the time at $d = 4$.
- An observed time of 25 leaves a residual. A high correlation does not license extrapolation.

Exponentials, logarithms and modelling

■ Why does a decay model stay positive?

- A medicine concentration falls by the same percentage each hour. A constant subtraction would eventually predict a negative amount.
 - This lesson studies **half-life** : The time for a decaying quantity to fall to half its initial value.
-

■ The method


- For $y = Ae^{kt}$, k is a proportional rate.
 - Taking logs gives $\ln y = \ln A + kt$.
 - Logarithms require positive arguments, and $\log(x+y)$ is not $\log x + \log y$.
-

■ A worked case

$$y = Ae^{kt}, \quad \ln y = \ln A + kt$$

Example:

If $y = 80e^{(-0.2t)}$, $y = 40$ gives $e^{(-0.2t)} = 1/2$. Hence $t = \ln 2 / 0.2 \approx 3.466$. For $3^x = 20$, $x = \ln 20 / \ln 3$. Plotting $\ln y$ against t linearises this exponential model.



$y = 80e^{(-0.2x)}$ loses the same fraction, about 18%, in every unit of x

■ Where it goes wrong

- A fitted exponential is a model, not a guarantee.
- Specify the time units and range of use.
- A negative k describes decay; a negative starting amount usually contradicts the context.

Not true: For positive x and y , $\ln(x+y) \neq \ln x + \ln y$. The log of a sum has no simple law; the product rule is $\ln(xy) = \ln x + \ln y$.

■ Applying it

- Compare actual observations with the model.
 - Systematic departures may indicate changing conditions.
 - In a report, explain what the rate and initial value mean, rather than giving a bare equation.
-

■ In your course

- Applications and Interpretation SL, the course first assessed in 2021.
- Every paper allows a calculator, but method marks still need the expression or equation written before its value.

Key:

- **Half-life:** The time for a decaying quantity to fall to half its initial value.
- For $y=Ae^{kt}$, k is a proportional rate.
- Watch for: A fitted exponential is a model, not a guarantee.

■ **Write from 2 to the power 7**

- Solve 2 to the power $x = 128$.
- The sum of $\ln 4$ and $\ln 7$ is $\ln 28$, not $\ln$ of the sum.

■ **Write from a half-life of 10 days**

- 200 g of a substance decays to 50 g in 20 days.
- Two halvings in 20 days, so the half-life is 10 days – count the halvings, not the subtraction.