

# Statistics and probability

International Baccalaureate · IB Diploma · Mathematics:  
Applications and Interpretation · SL

## Scope and prerequisites

---

This handout covers selected SL topics for the course first assessed in 2021; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.

## Data summaries, histograms and interpretation

---

### ■ Can one average tell the whole story?

- Two groups have the same median but different spread. One summary cannot describe both location and consistency.
  - This lesson studies **frequency density** : Frequency divided by class width, used as histogram height.
- 

### ■ The method


- Compare an appropriate average and spread in context.
  - A histogram uses area for frequency, so height=frequency/class width.
  - Grouped estimates assume representative values within intervals.
- 

### ■ A worked case

$$\bar{x} = \frac{\sum x_i}{n}, \quad \text{density} = \frac{\text{frequency}}{\text{class width}}$$

### Example:

A class from 10 to 20 with frequency 30 has density  $30/10=3$ . A class from 20 to 40 with frequency 20 has density  $20/20=1$ . Its wider bar must not be mistaken for a larger density. For values 2,4,4,6,9, the median is 4 and mean is 5.



*For 2, 4, 4, 6, 9: mean =  $25 \div 5 = 5$ , median = 4*

### ■ Where it goes wrong

- The tallest histogram bar need not contain the most observations.
- A grouped mean is an estimate.
- Correlation does not prove causation, and extrapolation extends beyond the observed range.

**Not true:** A histogram bar's height always equals its frequency, even with unequal class widths. With unequal class widths the height is frequency density; frequency is the area of the bar.

### ■ Applying it

- Choose a display that fits the data type.
- Give both a numerical comparison and what it means for the population; do not infer more precision than the sample supports.

### ■ In your course

- Applications and Interpretation SL, the course first assessed in 2021.
- Every paper allows a calculator, but method marks still need the expression or equation written before its value.

**Key:**

- **Frequency density:** Frequency divided by class width, used as histogram height.
- Compare an appropriate average and spread in context.
- Watch for: the tallest histogram bar need not contain the most observations.

### ■ Write from a mean of 8

- A class of width 3 has frequency 18.
- The values 4, 4, 7, 9 and 16 have a mean and a median. Bar height is not always the frequency.

## Probability, trees and conditional reasoning

---

### ■ What changes after the first draw?

- A bag has 3 red and 2 blue counters. Taking two without replacement changes the chance of the second colour.
  - This lesson studies **conditional probability** : The probability of an event after restricting the sample space to a stated condition.
- 

### ■ The method


- Multiply along a tree branch and add disjoint branches.
  - With replacement, the composition stays fixed.
  - Conditional probability is  $P(A \text{ given } B) = P(A \cap B) / P(B)$ , for  $P(B) > 0$ .
- 

### ■ A worked case

$$P(A | B) = \frac{P(A \cap B)}{P(B)}, \quad P(B) > 0$$

### Example:

Without replacement,  $P(\text{two red}) = 3/5 \times 2/4 = 3/10$ .  $P(\text{one of each}) = 3/5 \times 2/4 + 2/5 \times 3/4 = 3/5$ .  
 If  $P(A \cap B) = 0.12$  and  $P(B) = 0.3$ ,  $P(A \text{ given } B) = 0.4$ .



Without replacement:  $P(RR) = \frac{3}{5} \times \frac{2}{4} = 0.3$  and  $P(BB) = \frac{2}{5} \times \frac{1}{4} = 0.1$

### ■ Where it goes wrong

- Mutually exclusive means no overlap; independent means that knowing one event does not change the other's probability.
- Two disjoint events with positive probability are not independent.

**Not true:** Mutually exclusive events with positive probabilities must be independent. Mutually exclusive events with positive probabilities are dependent: if one happens, the other cannot, so  $P(A \text{ and } B) = 0 \neq P(A)P(B)$ .

### ■ Applying it

- A two-way table makes the restricted denominator visible.
- Before using a product  $P(A)P(B)$ , justify independence from the context or the supplied information.

### ■ In your course

- Applications and Interpretation SL, the course first assessed in 2021.
- Every paper allows a calculator, but method marks still need the expression or equation written before its value.

**Key:**

- **Conditional probability:** The probability of an event after restricting the sample space to a stated condition.
- Multiply along a tree branch and add disjoint branches.
- Watch for: mutually exclusive means no overlap; independent means that knowing one event does not change the other's probability.

### ■ Write from a conditional probability of 0.70

- $P(A \text{ and } B)$  is 0.28 and  $P(B)$  is 0.40.
- Find  $P(A \text{ given } B)$ . Mutually exclusive events that can happen are not independent.

## Binomial, normal and Poisson models

---

### ■ What makes a count predictable?

- A quality inspector counts defective items. The number is random, but a model can describe its likely range.
  - This lesson studies **expected value** : The probability-weighted mean of a random variable.
- 

### ■ The method


- A binomial model needs fixed  $n$ , independent trials, two outcomes and constant  $p$ .
  - $E(X)=np$  and  $\text{Var}(X)=np(1-p)$ .
  - For a normal model use  $z=(x-\mu)/\sigma$ .
  - A Poisson model describes counts with a constant rate and appropriate independence assumptions.
- 

### ■ A worked case

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad X \sim B(n, p)$$

### Example:

For  $X$  binomial(5,0.2),  $P(X=0)=0.8^5=0.32768$ ,  $E(X)=1$  and  $\text{Var}(X)=0.8$ . For a normal quantity with  $\mu=100, \sigma=15$ , the value 130 has  $z=2$ . A Poisson mean of 3 per hour gives mean 6 over two hours.



$X \sim B(5, 0.2)$ :  $P(X = 1) \approx 0.41$  is the largest bar;  $P(X = 5)$  is tiny

### ■ Where it goes wrong

- Not every count is binomial: changing  $p$  or dependence can invalidate it.
- For a continuous variable, the probability of one exact value is zero.
- Continuity correction matters when approximating a discrete distribution by a normal one.

**Not true:** Every count of successes has a binomial distribution regardless of dependence. The binomial model needs independent trials with a fixed probability; dependent trials break it.

### ■ Applying it

- Write the event as an inequality before using calculator distribution functions.
- Distinguish  $P(X < k)$ ,  $P(X \leq k)$  and a tail complement.
- State assumptions in context.

### ■ In your course

- Applications and Interpretation SL, the course first assessed in 2021.

### Key:

- **Expected value:** The probability-weighted mean of a random variable.
- A binomial model needs fixed  $n$ , independent trials, two outcomes and constant  $p$ .

- Watch for: not every count is binomial: changing  $p$  or dependence can invalidate it.
- **Write from a z-score of 4**
- A value 90 comes from a distribution with mean 70 and standard deviation 5.
  - Find the z-score. A binomial count needs independent trials.

## Hypothesis testing and contextual conclusions

---

■ **Could chance explain the result?**

- A factory claims that only 10% of items are defective. A sample contains more defects, but chance alone may explain some difference.
  - This lesson studies **significance level** : The chosen probability threshold for rejecting a null hypothesis.
- 

■ **The method**


- State  $H_0$  and  $H_1$  in population parameters before inspecting the outcome.
  - Calculate the appropriate tail probability under  $H_0$ .
  - Reject  $H_0$  when the evidence meets the specified significance rule; otherwise say there is insufficient evidence.
- 

■ **A worked case**

$$H_0 : p = p_0, \quad H_1 : p > p_0$$

**Example:**

For  $H_0:p=0.1$  and  $H_1:p>0.1$  with  $n=10$ , observing at least 3 defects has probability  $1-(0.9^{10}+10\times0.1\times0.9^9+45\times0.1^2\times0.9^8)\approx0.070191$ . At 5%, this is insufficient evidence that the defect rate exceeds 10%.



*Under  $B(10, 0.1)$ ,  $P(X \geq 3) \approx 0.07$ : the shaded upper tail is the evidence region for the test*

---

### ■ Where it goes wrong

- Failing to reject  $H_0$  is not proof that  $H_0$  is true.
- Choose the tail from  $H_1$ , not from whichever tail gives a small result.
- Statistical significance does not measure the practical size of an effect.

**Not true:** Failing to reject a null hypothesis proves it is true. Not rejecting the null hypothesis only means there is not enough evidence against it.

---

### ■ Applying it

- Finish with a sentence about the population and the original claim.
  - State the model's assumptions and consider whether the sampling procedure supports them.
- 

### ■ In your course

- Applications and Interpretation SL, the course first assessed in 2021.
- Every paper allows a calculator, but method marks still need the expression or equation written before its value.

**Key:**

- **Significance level:** The chosen probability threshold for rejecting a null hypothesis.
- State  $H_0$  and  $H_1$  in population parameters before inspecting the outcome.
- Watch for: failing to reject  $H_0$  is not proof that  $H_0$  is true.

■ **Write from a p-value of 0.08**

- A one-tailed p-value is 0.08 and the significance level is 5 percent.
- Decide whether to reject the null hypothesis. Failing to reject does not prove it true.

■ **Write from a p-value of 0.03**

- A one-tailed p-value is 0.03 against a 5 percent significance level.
- 0.03 is below 0.05, so reject  $H_0$  – the result is significant at the chosen level.

■ **Write from a two-tailed p-value of 0.08**

- A two-tailed p-value of 0.08 is tested at the 10 percent level.
- Reject at 10 percent; the identical result at 5 percent would not – the threshold is chosen BEFORE the data are seen.