

20.2 Structure of the atom (full E.1, HL)


Name: _____ Class: _____ Date: _____ Total: 30 marks

KEY WORDS nucleon number • proton number • emission spectrum
• Bohr model • distance of closest approach

Objective

Practise the full guide E.1 toolkit including AHL: the Geiger–Marsden–Rutherford experiment and the discovery of the nucleus, nuclear notation ${}^A_ZX$, emission and absorption spectra as evidence for discrete energy levels, photon emission and absorption in transitions with $E = hf$, and spectra as fingerprints of chemical composition; at higher level also $R = R_0A^{1/3}$ with nuclear density, the distance of closest approach, and the Bohr levels $E_n = -13.6/n^2$ eV with $mvr = nh/2\pi$. Original practice; not an official IB paper.

1 Key knowledge and worked examples



■ The E.1 rule set (HL)

$$E = hf \quad {}^A_ZX \quad R = R_0A^{1/3} \quad E_n = -\frac{13.6}{n^2} \text{ eV} \quad mvr = \frac{nh}{2\pi}$$

Alpha scattering: most particles pass through (empty atom), few deflect far (tiny dense positive nucleus). Spectral lines correspond one-to-one with transitions between discrete levels; the set of lines identifies the element.

■ Worked example: reading a line

Levels at -1.5 eV and -3.4 eV: a downward jump emits $E = 1.9$ eV $= 3.0 \times 10^{-19}$ J, so $f = E/h = (3.0 \times 10^{-19})/(6.6 \times 10^{-34}) = 4.6 \times 10^{14}$ Hz. The atom absorbs exactly this frequency climbing up – emission and absorption lines match. Two habits: convert eV to joules before using $E = hf$, and remember each line is one gap, not one level.

2 Practice

2.1 State what the Geiger–Marsden–Rutherford experiment revealed about the structure of the atom, citing two observations. [3]

2.2 Write the meaning of each symbol in nuclear notation ${}^A_ZX$. [2]

2.3 Why do emission and absorption spectra consist of sharp lines rather than continuous bands? [2]

2.4 State one use of spectral analysis that identifies chemical composition at astronomical distances. [1]

2.5 A 4.0 MeV alpha particle heads straight at a gold nucleus ($Z = 79$). Calculate its distance of closest approach. [3]

3 Exam-style questions

3.1 An atom has energy levels at -1.5 eV , -3.4 eV and -13.6 eV . Calculate the frequency of the photon emitted in the $-3.4 \rightarrow -1.5\text{ eV}$ transition, and the energy (in eV) of the photon that ionizes the atom from the -1.5 eV level. [4]

3.2 Explain why almost all alpha particles passed straight through the gold foil while about 1 in 8000 bounced back, and what each observation implies. [4]

3.3 A gas of element X shows absorption lines at exactly the same frequencies as its emission lines. Explain how this identifies X in the atmosphere of a distant star. [3]

3.4 The nuclide ${}_{92}^{235}\text{U}$ is used in reactors. State its proton number, nucleon number, and the number of neutrons in its nucleus. [3]

3.5 In the Bohr model of hydrogen, calculate the energy of the $n = 3$ level, the energy and wavelength of the photon emitted in the $3 \rightarrow 2$ transition ($hc = 1.99 \times 10^{-25}\text{ J m}$), and state the assumption about angular momentum that produces these levels. [5]
