

Wave model

International Baccalaureate · IB Diploma · Physics · HL

Scope and prerequisites

This handout covers selected HL topics for the course first assessed in 2025; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Waves, measurement and refraction

■ What would explain this observation?

- A wave can carry energy across water while a floating marker mainly oscillates. Energy transfer and bulk transfer of matter are different.
-

■ Build the model

- **Frequency** is oscillations per time; **wavelength** is distance between successive points in phase.
- Wave speed equals frequency multiplied by wavelength.
- Transverse oscillations are perpendicular to propagation.
- wavelength: Distance between successive points in phase.
- frequency: Number of oscillations per unit time.



One wavelength is the length of one complete cycle; $v = f\lambda$

■ Choose evidence that can test it

- At a boundary, frequency stays fixed by the source.
- A change of speed changes wavelength.
- Refraction follows from speed differences; angles are measured from the normal.

In the lab: Measure several wavelengths and divide to reduce fractional reading uncertainty. Use a ray box with a normal drawn at the boundary. Keep the beam away from eyes and record incident and refracted angles clearly.

■ Work from known quantities

- Known: frequency 5 Hz and wavelength 0.40 m.
- Use $v = f\lambda$. $v = 5 \times 0.40 = 2.0$ metres per second.
- If speed falls to 1.5 metres per second at the same frequency, wavelength = $v/f = 1.5/5 = 0.30$ m.

Your turn: Frequency is 8 Hz and wavelength is 0.25 m. Find wave speed.

■ Check the conclusion and its limits

- The wave frequency does not change simply because the medium changes.
- A longitudinal wave has oscillations parallel to propagation.

Not true: Wave frequency always changes when a wave enters a new medium. Frequency is fixed by the source, so it stays the same across a boundary. What changes is speed, and therefore wavelength through $v = f \times \text{wavelength}$.

Key:

- Frequency is oscillations per time; wavelength is distance between successive points in phase.
- At a boundary, frequency stays fixed by the source.
- Watch for: the wave frequency does not change simply because the medium changes.

■ Write from a wave speed of 7.2 metres per second

- Frequency is 9 Hz and wavelength is 0.80 m.
- Find the wave speed. If the speed becomes 5.4 metres per second at a boundary, find the new wavelength. Frequency stays the same.

■ **Write from a snell check**

- Light passes from air into glass ($n = 1.5$) at 30° to the normal.
- $\sin \theta_{\text{glass}} = \sin 30^\circ \div 1.5 = 0.333$, so $\theta_{\text{glass}} \approx 19.5^\circ$ – toward the normal in the slower medium, as it must be.

■ **Write from a critical angle**

- Light travels from glass ($n = 1.5$) toward air.
- $\sin \theta_c = 1 \div 1.5 = 0.667$, so $\theta_c \approx 41.8^\circ$ – beyond it, total internal reflection: the fibre-optic condition.