

Fusion and stars

International Baccalaureate · IB Diploma · Physics · SL

Scope and prerequisites

This handout covers selected SL topics for the course first assessed in 2025; it is not the whole syllabus.

Prerequisites: read the stated quantities and units, and use arithmetic and the model conditions below.

The examples are original and fictional: they are not real measurements or real student investigations.


Stars, radiation and scale

■ What would explain this observation?

- Two stars can have the same apparent brightness while having different luminosities. Distance changes the **flux** reaching an observer.
-

■ Build the model

- **Luminosity** is total power emitted; flux is power received per area.
- A stellar spectrum carries information about surface temperature and composition.
- Fusion transfers energy as light nuclei combine.
- luminosity: Total emitted power.
- flux: Power received per unit area.



Inverse square: the same power spread over $4\pi d^2$, so twice as far gives a quarter of the flux

■ Choose evidence that can test it

- For isotropic emission without absorption, flux follows an inverse-square relationship with distance.
- Observed brightness alone therefore cannot establish luminosity.

In the lab: Keep distance units consistent, identify which quantities are intrinsic to the star, and distinguish observational evidence from a model of stellar evolution. Do not confuse a red giant stage with every possible final remnant.

■ Work from known quantities

- Known: distance doubles while luminosity remains fixed.
- Use F proportional to $1/d^2$.
- Flux ratio = $1/2^2 = 1/4$.
- A flux of 12 units becomes 3 units.
- This assumes no change in absorption or source output.

Your turn: A source gives flux 18 units. Find flux at three times the distance.

■ Check the conclusion and its limits

- The Sun is not expected to become a supernova.
- Redshift can support cosmological expansion; it does not mean every nearby object must move away from every observer.

Not true: Apparent brightness alone always determines a star luminosity. Apparent brightness depends on distance as well as luminosity ($F = L/4\pi d^2$). A dim-looking star may be very luminous but far away, so brightness alone cannot fix luminosity without a distance.

Key:

- Luminosity is total power emitted; flux is power received per area.
- For isotropic emission without absorption, flux follows an inverse-square relationship with distance.
- Watch for: the Sun is not expected to become a supernova.

■ Write from a flux of 8 units

- A source gives flux 32 units.
- Find the flux at twice the distance. Apparent brightness alone does not give luminosity.

■ **Write from a distance tripled**

- A source gives flux 45 units at distance d .
- At $3d$: $45 \div 9 = 5$ units – the inverse-square law, one more time.

■ **Write from an apparent-brightness trap**

- Sirius outshines Betelgeuse in our sky, yet Betelgeuse emits far more light.
- Apparent brightness folds in distance ($F = L \div 4\pi d^2$) – “brightest in the sky” is a claim about d as much as L .

■ **Write from a luminosity comparison**

- Star A delivers twice the Sun’s power at twice the Sun’s distance.
- Its flux here = $2 \div 2^2 =$ half the Sun’s – luminosity is emission; flux is what ARRIVES, divided by $4\pi d^2$.