

4.2 Thermal energy transfers (full B.1)


Name: _____ Class: _____ Date: _____ Total: 25 marks

KEY WORDS internal energy • specific latent heat • emissivity

Objective

Practise the full guide B.1 toolkit: Kelvin temperature and mean kinetic energy ($E_k = \frac{3}{2} k_B T$), $Q = mc\Delta T$ and $Q = mL$, the conduction rate $\frac{\Delta Q}{\Delta t} = \frac{kA\Delta T}{\Delta x}$, Stefan–Boltzmann $L = \sigma AT^4$ with apparent brightness $b = \frac{L}{4\pi d^2}$, Wien’s $\lambda_{\text{max}} T = 2.9 \times 10^{-3} \text{ m} \cdot \text{K}$, and emissivity. Original practice; not an official IB paper.

1 Key knowledge and worked examples



■ The B.1 equation set

$E_k = \frac{3}{2} k_B T$ $Q = mc\Delta T$ $Q = mL$ $\frac{\Delta Q}{\Delta t} = \frac{kA\Delta T}{\Delta x}$

$L = \sigma AT^4$ $b = \frac{L}{4\pi d^2}$ $\lambda_{\text{max}} T = 2.9 \times 10^{-3} \text{ m} \cdot \text{K}$ $\text{emissivity} = \frac{\text{power per unit area}}{\sigma T^4}$

Worked check. A star radiates like a black body of radius $7.0 \times 10^8 \text{ m}$ at 5800 K :
 $L = 5.7 \times 10^{-8} \times 4\pi(7.0 \times 10^8)^2 \times 5800^4 \approx 4 \times 10^{26} \text{ W}$.

2 Practice

2.1 Convert 27°C to kelvin, and a change of 45 K to Celsius degrees. [2]

2.2 Calculate the mean kinetic energy of a gas molecule at 300 K ($k_B = 1.38 \times 10^{-23} \text{ J/K}$). [2]

2.3 0.50 kg of water cools from 80°C to 30°C ($c = 4186 \text{ J/(kg} \cdot \text{K)}$). Calculate Q . [2]

2.4 A metal rod ($k = 50$, $A = 2.0 \times 10^{-4} \text{ m}^2$, length 0.40 m) has ends 100 K apart. Calculate the conduction rate. [3]

2.5 A black body’s spectrum peaks at 970 nm . Calculate its temperature. [2]

3 Exam-style questions

3.1 A 1.2 kW heater warms 2.0 kg of ice at 0°C . Calculate the energy to melt the ice ($L_f = 3.34 \times 10^5 \text{ J/kg}$), the further energy to reach 25°C , and the minimum total heating time. [5]

3.2 A planet orbits a star of luminosity 4.0×10^{26} W at a distance of 1.5×10^{11} m. Calculate the apparent brightness, and the power incident on the planet's cross-section (πR^2 , $R = 6.4 \times 10^6$ m). [5]

3.3 A radiator surface (2.0 m^2 , 340 K) has emissivity 0.80 . Calculate the power radiated, and compare it with a black body at the same temperature. [4]
