

8.2 Simple harmonic motion (full C.1, SL)

Name: _____ Class: _____ Date: _____

Total: 22 marks

KEY WORDS simple harmonic motion
• equilibrium position


• restoring force

• amplitude

Objective

Practise the full guide C.1 toolkit at standard level: the defining condition $a = -\omega^2 x$ and the meaning of its minus sign, $T = 1/f = 2\pi/\omega$, the mass-spring period $T = 2\pi\sqrt{m/k}$, the pendulum period $T = 2\pi\sqrt{l/g}$, and qualitative energy exchange through a cycle. Original practice; not an official IB paper.

1 Key knowledge and worked examples



■ The C.1 equation set (SL)

$$a = -\omega^2 x \quad T = \frac{1}{f} = \frac{2\pi}{\omega} \quad T = 2\pi\sqrt{\frac{m}{k}} \quad T = 2\pi\sqrt{\frac{l}{g}}$$

Restoring condition: acceleration proportional to displacement and directed back toward equilibrium. Energy: total constant, kinetic greatest at equilibrium, potential greatest at the extremes.

■ Worked example: rating a spring–mass oscillator

A 0.50 kg mass hangs on a spring of stiffness 20 N/m and is pulled down and released. Period $T = 2\pi\sqrt{0.50/20} = 0.99$ s, so $f \approx 1.0$ Hz and $\omega = 6.3$ rad/s. With amplitude 0.050 m the maximum acceleration is $\omega^2 x_0 = 158 \times 0.050 = 7.9$ m/s², at the extremes where the mass momentarily stops. Two habits: check that the restoring condition holds before using these equations, and place maximum speed at equilibrium, never at the extremes.

2 Practice

2.1 State the defining equation of simple harmonic motion and explain the significance of its minus sign. [2]

2.2 An oscillator has frequency 2.5 Hz. Calculate its period and angular frequency. [2]

2.3 A 0.20 kg mass on a spring of stiffness 5.0 N/m oscillates. Calculate the period. [2]

2.4 Describe the energy changes over one complete cycle of a simple pendulum, from release at maximum displacement. [3]

3 Exam-style questions

3.1 A simple pendulum has length 2.5 m ($g = 9.8 \text{ m/s}^2$). Calculate its period, and calculate the length that would double this period. [4]

3.2 A 0.30 kg mass on a spring oscillates with amplitude 0.040 m and period 0.25 s. Calculate the angular frequency and the maximum acceleration, and state where in the cycle the speed is greatest. [4]

3.3 A student times 20 oscillations of a mass–spring system as 8.8 s, with mass 0.40 kg. Calculate the period and the spring stiffness, and state one assumption the calculation makes about the system. [5]
