5.1 Motion, Newton's laws and impulse

Name: _____ Class: _____ Date: _____ **Total: 30 marks**

KEY WORDS velocity • acceleration • ground reaction force • momentum • impulse • Newton's third law

Objective

Build the skills to answer exam questions on describing and explaining motion: velocity and acceleration, motion graphs, **Newton's laws** , momentum and **impulse** .

You must be able to:

- use distance, displacement, speed, velocity and acceleration with units
- read motion graphs: gradient and area
- apply Newton's three laws to sporting movements
- calculate momentum and impulse, and explain why longer contact times reduce force

1 Key knowledge and worked examples

Motion and Newton's laws

Scalars have size only (distance, speed, mass); **vectors** have size and direction (displacement, velocity, force).

$$v = \frac{\text{displacement}}{t} \qquad a = \frac{\Delta v}{t} \qquad p = mv \qquad \text{impulse} = F\,\Delta t = \Delta p$$

Graphs: on a displacement–time graph, the **gradient** is velocity. On a velocity–time graph, the gradient is **acceleration** and the **area** under the line is **displacement**. On a force–time graph, the area is **impulse**.

law	statement	sport example
first (inertia)	a body stays at rest or moves at constant velocity unless a resultant force acts	a ball keeps rolling until friction slows it
second	$F = ma$: resultant force = mass × acceleration	a bigger push gives a bigger acceleration
third	every action force has an equal and opposite reaction force	the sprinter pushes back on the blocks; the blocks push the sprinter forward (**ground reaction force**)

Impulse: the same change in momentum can come from a large force for a short time or a smaller force for a longer time. Bending the knees on landing or "giving" with the hands when catching makes the contact time longer, so the force is smaller and injury risk lower.

AHL: in running each foot contact gives a **braking** impulse then a **propulsive** impulse; the runner speeds up when the propulsive impulse is larger. From a jump's net impulse, take-off velocity $v = \text{impulse} \div m$, and height $h = v^2 \div 2g$ with $g = 9.81$ m s^{-2}.

Answer shapes: formula → substitute with units → answer with units and direction if it is a vector.


2 Practice

2.1 Distinguish between speed and velocity. [2]




exercises.lol exercises.com.cn


2.2 A 70 kg sprinter goes from rest to 10 m s^{-1} in 2.5 s. Calculate her mean acceleration and the mean resultant force on her. [3]



2.3 Use Newton's third law to explain how a sprinter is pushed forward out of the blocks. [3]




2.4 The force–time graph shows a push-off lasting 0.2 s. The force rises from 0 to 1500 N and falls back to 0 in a triangle shape. The athlete has a mass of 75 kg. Calculate the impulse and the change in velocity. [3]



exercises.lol exercises.com.cn

3 Exam-style questions


3.1 The velocity–time graph shows the first 6 s of a 100 m race.



(a) Describe the motion of the sprinter. [2]

(b) Calculate the displacement in the first 2 s. [2]

3.2 A 40 kg gymnast lands at 5 m s^{-1}. With stiff legs she stops in 0.1 s; with bent knees she stops in 0.4 s.




(a) Calculate her momentum just before landing. [1]

(b) Calculate the mean stopping force in each landing (ignore her weight). [3]

(c) Explain why bending the knees reduces the risk of injury. [2]


3.3 The graph shows the horizontal ground reaction force during one foot contact in running.



(a) Explain why the first part of the contact slows the runner. [2]

(b) Explain how the runner can speed up over several steps. [2]

3.4 In a vertical jump on a force plate, the net upward impulse is 180 N s. The jumper's mass is 60 kg.



(a) Calculate the take-off velocity. [2]

(b) Calculate the jump height ($g = 9.81$ m s^{-2}). [2]

(c) Suggest **one** reason why a jump-and-reach test may give a different height. [1]

exercises.lol exercises.com.cn