

E6.5 Non-right-angled triangles

Name: _____ Class: _____ Date: _____

Total: 16 marks

KEY WORDS sine rule 正弦定理 • cosine rule 余弦定理 • included angle 夹角
• ambiguous case 两解情形 • obtuse 钝角

Objective


Build the skills to answer Extended exam questions on **non-right-angled triangles** 非直角三角形 — choosing between the **sine rule** 正弦定理 and the **cosine rule** 余弦定理, and using the **area formula** $\frac{1}{2}ab \sin C$ 面积公式.

You must be able to:

- choose the correct rule from what the question gives you
- use the sine rule to find a side or an angle
- use the cosine rule to find a third side, and use $\frac{1}{2}ab \sin C$ for an area

1 Worked examples

■ Which rule? Count what you are given



The angle in the cosine rule sits between the two known sides

- **Sine rule:** $\frac{a}{\sin A} = \frac{b}{\sin B}$. Use it when you have a side with its OPPOSITE angle, plus one more piece.
- **Cosine rule:** $a^2 = b^2 + c^2 - 2bc \cos A$. Use it for two sides and the angle BETWEEN them, or for all three sides.
- **Area:** $\frac{1}{2}ab \sin C$, where C is the angle between the two sides a and b .
- **The ambiguous case:** when the sine rule gives an angle, 180° minus that angle also has the same sine. Check which fits the triangle.

Answer shapes (marked by **method marks** — M1 for a correct method, A1 for the correct answer): quoting the rule with the numbers substituted is the method mark, and the rearranged line is a second one.

2 Practice

2.1 In triangle ABC , $a = 9$ cm, $A = 40^\circ$ and $B = 65^\circ$. Find b , correct to 3 significant figures. [3]

2.2 In triangle PQR , $p = 7$ cm, $q = 9$ cm and $R = 52^\circ$. Find r , correct to 3 significant figures. [3]

2.3 Find the area of a triangle with sides 10 cm and 14 cm and an included angle of 38° , correct to 3 significant figures. [2]

3 Exam-style questions

3.1 In triangle ABC , $AB = 8$ cm, $BC = 11$ cm and angle $ABC = 73^\circ$.

(a) Find AC , correct to 3 significant figures.

[3]

(b) Find the area of the triangle, correct to 3 significant figures.

[2]

3.2 In triangle XYZ , $x = 6.2$ cm, $y = 9.4$ cm and angle $X = 35^\circ$. Find the acute value of angle Y , correct to 1 decimal place.

[3]
