

Exercise walk-through

13.5


Optimisation and learning

University Computer Science

Optimisation and learning: descend, sized and scaled

- 1 **Quadratic** $L(w) = (w-3)^2$: eta 0.1 walks 0.6, 1.08, 1.464, 1.771, 2.017; eta 1 oscillates 0, 6 forever; stable iff eta < 1.
- 2 **Line** On the least-squares line: two steps from origin with eta 0.02 reach (0.84, 0.25); a thousand close on (0.9, 1.3).
- 3 **Ravine** $L = (x_1-1)^2 + 100(x_2-1)^2$: one rate cannot serve both curvatures - standardise features.

Loss surface and updates



Three learning rates on the same bowl: crawl, damped, oscillation.

Exercises 2.1-2.3

2.1 For $L(w) = (w - 3)^2$ with $\eta = 0.1$ from $w_0 = 4$: give w_1, w_2 . [3]

2.2 What is the stability bound on η here? [3]

2.3 What shape does $L = (x_1 - 1)^2 + 100(x_2 - 1)^2$ have? [3]

Exercises 2.1-2.3

2.1 For $L(w) = (w - 3)^2$ with $\eta = 0.1$ from $w_0 = 4$: give w_1, w_2 . [3]

2.1

- $w_1 = 4 - 0.2(1) = 3.8$; $w_2 = 3.8 - 0.2(0.8) = 3.64$ ✓.

2.2 What is the stability bound on η here? [3]

2.2

- $|1 - 2\eta| < 1 \iff 0 < \eta < 1$ ✓.

2.3 What shape does $L = (x_1 - 1)^2 + 100(x_2 - 1)^2$ have? [3]

2.3

- A ravine: 100x curvature mismatch between the axes ✓.

Exercise 3.1

3.1 Run gradient descent on $L(w) = (w + 2)^2$ (note the sign) from $w_0 = 1$ with $\eta = 0.25$ for four steps.

Give the update rule specialised to this loss. List every iterate (exact fractions or 4 decimals). State the limit and whether the steps overshoot. Give the iterate sequence for $\eta = 0.75$ and state its qualitative difference. [8]

Exercise 3.1

3.1 Run gradient descent on $L(w) = (w + 2)^2$ (note the sign) from $w_0 = 1$ with $\eta = 0.25$ for four steps.

Give the update rule specialised to this loss. List every iterate (exact fractions or 4 decimals). State the limit and whether the steps overshoot. Give the iterate sequence for $\eta = 0.75$ and state its qualitative difference. [8]

3.1

- Update: $w_{t+1} = w_t - 0.5(w_t + 2) = 0.5w_t - 1$ ✓.
- Iterates: $1 \rightarrow -0.5 \rightarrow -1.25 \rightarrow -1.625 \rightarrow -1.8125$ ✓.
- Limit -2 (the minimum); errors shrink monotonically, no overshoot ✓.
- With $\eta = 0.75$: $w_{t+1} = w_t - 1.5(w_t + 2) = -0.5w_t - 3$: $1 \rightarrow -3.5 \rightarrow -1.25 \rightarrow -2.375 \rightarrow -1.8125$ – alternating sign of the error, damped oscillation ✓.

Exercise 3.2

3.2 Explain, with numbers, why feature scaling changes the usable learning rate.

For $L = (x_1 - 1)^2 + 100(x_2 - 1)^2$ from $(0, 0)$: state the curvature along each axis and the stability bound on η for a single shared rate. Give the number of steps needed to reduce each coordinate's error by half under that bound. State what standardising the second feature does to the bound and to the descent. [7]

Exercise 3.2

3.2 Explain, with numbers, why feature scaling changes the usable learning rate.

For $L = (x_1 - 1)^2 + 100(x_2 - 1)^2$ from $(0, 0)$: state the curvature along each axis and the stability bound on η for a single shared rate. Give the number of steps needed to reduce each coordinate's error by half under that bound. State what standardising the second feature does to the bound and to the descent. [7]

3.2

- Curvatures: 2 and 200 (the loss's second derivatives) ✓.
- Shared-rate stability: $\eta < 2/200 = 0.01$ (the tighter axis rules) ✓.
- Halving the error per coordinate: the slow axis contracts at $|1 - 2\eta| = 0.98$ per step – about 34 steps to halve ✓.
- Standardising the second feature equalises curvature at 2: the bound becomes $\eta < 1$ and one rate serves both axes ✓.