

Discrete Mathematics

Curated computer science study track (not a degree, no licensure claim). Original explanations and practice only, following the reviewed course handout.

From counting on fingers to counting anything

- 2.1-2.3 the proof engine (logic, sets, induction); 2.4-2.6 the counting engine.
- Every proof names its strategy; every identity is checked on a small example first.

Logic and proofs: translate, then choose the shape

- 1 Equivalences** $p \rightarrow q \equiv \neg p \vee q$; $p \rightarrow q \equiv \neg q \rightarrow \neg p$; $\neg(p \wedge q) \equiv \neg p \vee \neg q$;
negation swaps $\forall$ with $\exists$.
- 2 Shapes** Direct (assume p , derive q) · **contraposition** (prove $\neg q \rightarrow \neg p$) ·
contradiction (assume $p \wedge \neg q$, absurdity).
- 3 Named** n^2 even $\rightarrow$ n even, by contraposition: $n = 2k+1$ gives $n^2 = 2(2k^2+2k)+1$, odd.

A truth table and its proof tree

the table checks

p	q	$p \rightarrow q$	$\neg q \rightarrow \neg p$
T	T	T	T
T	F	F	F
F	T	T	T
F	F	T	T

matching columns prove:

$p \rightarrow q \equiv \neg q \rightarrow \neg p$ (contraposition)

DIRECT

assume p, derive q


CONTRAPOSITION

prove $\neg q \rightarrow \neg p$

CONTRADICTION

assume $p \wedge \neg q$, absurdity

the tree structures




The table checks; the tree structures — three proof shapes choose your data.

Sets, functions and relations: arrows and matrices


- 1 **Sets** Union, intersection, difference, complement; the **power set** of A pairs in/out bits - one bit each.
- 2 **Functions** $f(n) = 2n$ **injective** not **surjective**; $\lfloor n/2 \rfloor$ surjective not injective; $-n$ bijective (own inverse).
- 3 **Relations** Reflexive = diagonal 1s; symmetric = matrix = transpose; transitive = check length-2 paths.

Set operations, arrow diagrams and relation properties


INJECTIVE



SURJECTIVE



BIJECTIVE



set algebra

$A \cup B$, $A \cap B$, $A - B$, complement
 power set: a bit string per subset
 $g \circ f$ reads right to left

relations as matrices


reflexive: diagonal all 1
 symmetric: matrix = its transpose
 transitive: check paths of length 2

Arrows classify the function; the matrix reads the relation at a glance.

Induction and recursion: the ladder

- 1 Ladder** Base $P(n_0)$ by hand; step: assume $P(k)$, derive $P(k+1)$ - induction is chained modus ponens.
- 2 Sum** $\sum_1^n i = n(n+1)/2$: base $1 = 1 \cdot 2/2$; step $k(k+1)/2 + (k+1) = (k+1)(k+2)/2$.
- 3 Faults** No base (ladder on air); hypothesis unused; step silently needs $k \geq 2$ - the horses proof fails at $k+1 = 2$.

The induction ladder: base, hypothesis, step



$\Sigma i = n(n+1)/2$, proved

base: $1 = 1 \cdot 2/2 \checkmark$

step: assume $\Sigma_{i=1}^k i = k(k+1)/2$
 then $\Sigma_{i=1}^{k+1} i = k(k+1)/2 + (k+1)$
 $= (k+1)(k+2)/2 \checkmark$

the hypothesis is USED,
 the algebra is SHOWN

the three faults

no base: a ladder on air
 hypothesis never used
 step needs $k \geq 2$ (all horses)


Base plants the ladder; the step climbs one rung.

Counting and combinatorics: two rules, two formulas

- 1 Rules** Disjoint alternatives ADD; sequential choices MULTIPLY - 4 routes $\times$ 3 vehicles = 12.
- 2 Formulas** $P(n,r) = n!/(n-r)!$ ordered; $C(n,r) = n!/(r!(n-r)!)$ unordered - the $r!$ is the whole difference.
- 3 IE** $|A \cup B \cup C| = \text{singles} - \text{pairs} + \text{triple}$: $45+35+30 - 12-10-8 + 5 = 85$; team of 3 from 8 with a leader: 168 both ways.

Counting trees: ordered vs unordered

the tree: ordered vs unordered



ordered pairs ($ab \neq ba$): $3 \times 2 = 6$

unordered $\{a,b\} = \{b,a\}$: $6 \div 2! = 3$

sequential choices MULTIPLY;
disjoint alternatives ADD

the two formulas

$P(n, r) = n!/(n-r)!$ ordered

$C(n, r) = n!/(r!(n-r)!)$ unordered

same choices; combinations divide by the $r!$ arrangements of each set

inclusion-exclusion

$|A \cup B \cup C| = \text{singles} - \text{pairs} + \text{triple}$

$45 + 35 + 30 - 12 - 10 - 8 + 5 = 85$

in none of the three: 15


Branches multiply, alternatives add, overcounts subtract.

Graphs and trees: one graph, three representations

- 1 **Represent Adjacency matrix** n^2 space with $O(1)$ edge test; **adjacency list** $O(|V|+|E|)$ - density decides.
- 2 **Handshake** Every edge adds 2: $\sum \deg = 2|E|$ - drawn graph $3+2+2+2+3 = 12 = 2 \cdot 6$; odd-degree count always even.
- 3 **Trees** Connected + acyclic $\rightarrow |E| = |V| - 1$; 12-vertex tree, 5 leaves $\rightarrow$ internal degree sum $22 - 5 = 17$.

Small explicit graphs with adjacency structures

the picture (5 vertices, 6 edges)



$$\sum \text{deg} = 3+2+2+2+3 = 12 = 2 \cdot 6 \checkmark$$

adjacency matrix

0	1	0	1	1
1	0	1	0	0
0	1	0	0	1
1	0	0	0	1
1	0	1	1	0

n^2 space, $O(1)$ edge test

adjacency list

1: 2 4 5
2: 1 3
3: 2 5
4: 1 5
5: 1 3 4

$O(|V|+|E|)$: sparse graphs

tree: connected + acyclic, so $|E| = |V| - 1$

12 vertices, 5 leaves: internal degree sum $22 - 5 = 17$

BFS: queue, level by level, shortest hops · DFS: stack, components

Three representations of one graph - and the degree sum checked by handshake.

Discrete probability: counting with weights

- 1 Sample** Two dice: 36 pairs; $P(\text{sum } 7) = 6/36 = 1/6$ - counting from 2.4 does the work.
- 2 Condition** $P(A|B) = P(A \cap B)/P(B)$ - the B-branch renormalised; independence multiplies $P(A)P(B)$.
- 3 Bayes** 1 % disease, $Se = Sp = 0.9$: $P(\text{dis}|+) = 0.009/(0.009+0.099) = 1/12 \approx 8.3\%$ - rare stays rare.

An outcome tree with conditional branches

the outcome tree (test for a 1 % disease)



$$\text{Bayes: } P(\text{dis} \mid +) = 0.009 / (0.009 + 0.099) \\ = 0.009 / 0.108 = 1/12 \approx 8.3 \%$$

the + branch keeps 0.108 of the world;
the diseased slice of it is 0.009

the rules

$P(E) = |E| / |\Omega|$
(equal outcomes: counting)

$P(A \mid B) = P(A \cap B) / P(B)$
(the B branch, renormalised)

Bayes:
 $P(A|B) = P(B|A) \cdot P(A) / P(B)$

independent:
 $P(A \cap B) = P(A) \cdot P(B)$

two dice: $P(\text{sum } 7) = 6/36 = 1/6$

The tree shows what conditioning removes: the branch, renormalised.

Check yourself

- **Power set** of a 3-element set...? (8 members.)
- $P(7,2)$ and $C(7,2)$...? (42 and 21.)
- Edges in a 12-vertex tree...? (11.)
- $P(\text{sum } 7)$ with two dice...? ($1/6$.)