

2.8 Vector calculus

Name: _____ Class: _____ Date: _____

Total: 21 marks

KEY WORDS gradient • divergence • curl • divergence theorem
• Stokes' theorem

Objective

Compute gradient, divergence and curl in Cartesian coordinates, use $E = -\nabla V$, and verify field results by a small-step line integral.

1 Key knowledge and worked examples


■ Three operators on fields

A vector field assigns a vector to each point; a scalar field a number. Three operators connect them:

- **Gradient** ∇V (scalar $\rightarrow$ vector): points up the steepest slope; the electric field is $E = -\nabla V$.
- **Divergence** $\nabla \cdot F$ (vector $\rightarrow$ scalar): net outflow per unit volume; for electric fields, the charge density.
- **Curl** $\nabla \times F$ (vector $\rightarrow$ vector): circulation per unit area; zero for electrostatics, the mechanism of induction where it is not.

For $F = (P, Q, R)$ in Cartesian coordinates: $\nabla \cdot F = \partial P/\partial x + \partial Q/\partial y + \partial R/\partial z$, and $\nabla V = (\partial V/\partial x, \partial V/\partial y, \partial V/\partial z)$.

Read a field at a glance: outflow means divergence, swirl means curl



■ Worked example: field from potential

$$V(x, y) = 3x^2y \text{ volts.}$$

- $\nabla V = (6xy, 3x^2)$; at $(1, 2)$: $\nabla V = (12, 3)$, so $E = -(12, 3)$ V/m.
- Check by a small step: move $\delta = (0.01, 0)$ from $(1, 2)$; V changes by $\approx \partial V / \partial x \cdot \delta x = 12 \times 0.01 = 0.12$ V, and $-E \cdot \delta = 12 \times 0.01 = 0.12$ V ✓. The field-times-step predicts the potential drop – that IS the gradient.

■ Reading the two theorems

The **divergence theorem** converts a volume integral of $\nabla \cdot F$ into a closed-surface flux; **Stokes' theorem** converts a surface integral of $\nabla \times F$ into a loop circulation. Maxwell's equations are these two statements with physics poured in – Gauss's law is the divergence theorem; Faraday's law is Stokes.

2 Practice

2.1 For $V = 5x^2$: compute ∇V and E , and evaluate both at $x = 2$. [3]

2.2 For $F = (3x, 2y, 0)$: compute $\nabla \cdot F$ at $(1, 1, 0)$ and say whether the point is a source or a sink. [3]

2.3 State the curl of any electrostatic field, and the physical law that changes this where magnetic fields vary. [2]

3 Exam-style questions

3.1 A potential is $V(x, y) = 10x - 5y$ volts.

Compute $\mathbf{E}$, state its magnitude and direction, and verify by computing the change in V along the step $(0.2, -0.1)$ from the origin two ways: directly from $\mathbf{V}$, and from $-\mathbf{E} \cdot \delta$. [6]

3.2 The field $F = (2xy, x^2, 0)$ is claimed to be conservative (curl-free).

Compute $\nabla \times F$ to verify the claim. Then compute the line integral of $\mathbf{F}$ along the straight path from $(0,0)$ to $(1,1)$ parameterised as (t, t) , and confirm the same value along the path (t, t^2) – path independence being the signature of a conservative field. [7]
