5.2 First law for closed systems

Name: _____ Class: _____ Date: _____ **Total: 14 marks**

KEY WORDS closed system • boundary work • adiabatic process


Objective

Keep the closed-system ledger: boundary work as area under the path, the first law with one sign convention, and the adiabatic relation.

1 Key knowledge and worked examples

■ The ledger

$\Delta U = Q - W$ (work by the system positive); $W = \int_1^2 p\,dV$ is the area under the path; reversible adiabatic: $pV^\gamma =$ constant with $\gamma = 1.4$ for air.



*The area under each path is that path's work

■ Worked example. Three paths, one pair of states

Isobaric to triple volume: $W = 200 \times 0.6 = 120$ kJ; isothermal: $W = 200 \times 0.3 \times \ln 3 = 65.9$ kJ ✓ - 54.1 kJ apart, exactly the area between the curves.

exercises.lol exercises.com.cn

2 Practice

Air (closed system) starts at $(300\ \text{kPa}, 0.2\ \text{m}^3)$ with $\gamma = 1.4$.

2.1 Air at 300 kPa, 0.2 m^3 expands isobarically to 0.6 m^3: the boundary work, and the heat transfer if $\Delta U = +84$ kJ (one decimal, kJ). [4]

2.2 The same gas, from 300 kPa, 0.2 m^3 to 0.6 m^3, connected isothermally instead: the boundary work (one decimal, kJ). [3]

2.3 The gas ($\gamma = 1.4$), starting again at 300 kPa, 0.2 m^3, is compressed adiabatically to 0.1 m^3: the final pressure (one decimal, kPa), the work (one decimal, kJ) and the change in internal energy (one decimal, kJ). [5]

2.4 One sentence each: why is boundary work path-dependent, and why does $\Delta U = mc_v\Delta T$ still hold on a non-constant-volume path for an ideal gas? [2]

exercises.lol exercises.com.cn