

10.6 Gastrointestinal, liver and renal pathology

Name: _____ Class: _____ Date: _____

Total: 24 marks

KEY WORDS cirrhosis • nephrotic syndrome

Objective

Trace portal hypertension; split the two renal syndromes.

1 Key knowledge and worked examples

■ The blocked vein

Cirrhosis (fibrosis + nodules) → portal hypertension → varices, splenomegaly (→ thrombocytopenia), ascites, caput medusae; failure adds encephalopathy (ammonia), jaundice, coagulopathy ✓.

■ Worked example: two barrier lesions

Nephritic: torn pores – haematuria with casts, hypertension, proteinuria <3.5 g/day.
Nephrotic: lost charge – ≥3.5 g/day, hypoalbuminaemia, oedema, hyperlipidaemia ✓. Panel: ALT/AST injury, ALP obstruction, albumin + INR synthesis.

2 Practice

2.1 Why does cirrhosis lower platelets? [3]

2.2 The nephrotic proteinuria line, with units? [3]

2.3 Which liver marker pair reports synthetic function? [3]

3 Exam-style questions

3.1 A fictional patient with cirrhosis: vomiting blood, distended abdomen, drowsy, bruising.

Localise the bleed (oesophageal varices – the portal collateral emergency) ✓. Explain the ascites (portal hydrostatic ↑ + low albumin oncotic ↓) and the bruising (vitamin-K factors and INR – synthetic failure) ✓. Attribute the drowsiness (ammonia → encephalopathy) ✓. [8]

3.2 Two fictional urines: A – Coca-Cola urine, red-cell casts, BP up, protein 1.2 g/day; B – frothy urine, protein 4.2 g/day, albumin 2.4 g/dL, periorbital oedema.

Classify A (nephritic – torn pores) and B (nephrotic – lost charge) with the quantitative line ✓. Explain why B oedemas (oncotic fall) while A is hypertensive (inflammatory renin and volume) ✓. [7]

Solutions

2.1

- Splenic sequestration – portal hypertension pools platelets in the enlarged spleen ✓.

2.2

- ≥ 3.5 g/day ✓.

2.3

- Albumin and INR ✓.

3.1

- Varices: the emergency portosystemic collateral ✓.
- Ascites: $\uparrow$ portal hydrostatic plus $\downarrow$ oncotic (low albumin) ✓.
- Bruising: synthetic failure – clotting factors, vitamin K, INR ✓.
- Drowsiness: ammonia crossing into brain – encephalopathy ✓.

3.2

- A: nephritic – blood with casts, hypertension, protein < 3.5 ✓.
- B: nephrotic – 4.2 g/day ≥ 3.5 , albumin 2.4 low, frothy urine ✓.
- B oedemas: albumin lost $\rightarrow$ oncotic falls $\rightarrow$ periorbital swelling ✓.
- A hypertensive: inflammation drives volume and renin ✓.